

# PAKISTAN JOURNAL OF HEALTH SCIENCES

(LAHORE)

https://thejas.com.pk/index.php/pjhs ISSN (E): 2790-9352, (P): 2790-9344 Volume 6, Issue 10 (October 2025)



## **Original Article**



Impact of Ghutka and Mawa Use on Oral Health in a Sub-Urban Population of Karachi

# Vijay Kumar Khatri¹, Hafsa Munir², Tooba Mahmud Gauhar³, Khalid Iqbal⁴, Devi Kumari⁵ and Roop Kumar⁵

- <sup>1</sup>Department of ENT, Al-Tibri Medical College and Hospital, Karachi, Pakistan
- <sup>2</sup>Department of Nephrology, Sindh Institute of Urology and Transplantation, Karachi, Pakistan
- <sup>3</sup>Department of General Surgery, Al-Tibri Medical College and Hospital, Karachi, Pakistan
- <sup>4</sup>Atia General Hospital, Malir, Karachi, Pakistan
- <sup>5</sup>Department of Anatomy, Al-Tibri Medical College and Hospital, Karachi, Pakistan
- <sup>6</sup>Jinnah Postgraduate Medical Center, Karachi, Pakistan

#### ARTICLE INFO

#### Keywords:

Ghutka, Mawa, Smokeless Tobacco, Oral Lesions, FNT

#### How to Cite:

Khatri, V. K., Munir, H., Gauhar, T. M., Iqbal, K., Kumari, D., & Kumar, R. (2025). Impact of Ghutka and Mawa Use on Oral Health in a Sub-Urban Population of Karachi: Impact of Ghutka and Mawa on Oral Health in Karachi. Pakistan Journal of Health Sciences, 6(10), 164–167. https://doi.org/10.54393/pjhs.v6i10.3358

## \*Corresponding Author:

Vijay Kumar Khatri Department of ENT, Al-Tibri Medical College and Hospital, Karachi, Pakistan doctor\_vijay@hotmail.co.uk

Received Date: 18<sup>th</sup> July, 2025 Revised Date: 17<sup>th</sup> October, 2025 Acceptance Date: 25<sup>th</sup> October, 2025 Published Date: 31<sup>st</sup> October, 2025

#### ABSTRACT

Ghutka and Mawa are smokeless tobacco products mostly used in South Asia. They have areca nut, tobacco, lime, catechu, paraffin wax, and flavoring materials. These mixtures are highly addictive and cancer-causing, related to oral submucous fibrosis, leukoplakia, and mouth cancers. Still, awareness of their harmful effects remains very low among users. Objectives: To observe oral findings and clinical patterns in Ghutka and Mawa users visiting ENT clinics. Methods: This cross-sectional research was done in Al-Tibri Medical College and Hospital, Karachi, from January to June 2025. One hundred patients who used Ghutka or Mawa for at least six months were selected by purposive sampling. Oral history and a detailed mouth examination were done. Data was entered and analyzed in SPSS using the Chi-square test, with a significance level p  $\leq$  0.05. **Results:** Out of 100 users, dental issues were most common (73%), then trismus (44%), pain (40%), and chewing trouble (21%). The majority were addicted for 6–20 years, while 11  $\,$ had more than 20 years of use. Ulcers were found in 58 Mawa and 53 Ghutka users; growths in 35 and 29 respectively. Longer use had more severe lesions, with malignant signs mostly after 10 years (p < 0.05). Conclusions: Ghutka and Mawa are strongly connected with ulcers, dental and jaw problems, and precancerous growths. Long-term use increases damage. Awareness, early detection, and strict public control are urgently required.

## INTRODUCTION

Ghutka and Mawa are smokeless tobacco products that are commonly used in the Indian subcontinent and also among some immigrant groups living abroad. They usually have areca nut, slaked lime, catechu, paraffin wax, tobacco, and different flavoring agents that can be sweet or spicy, mixed in a powder or small grain-like form [1-2]. When people chew them, the mixture quickly mixes with saliva and causes a reddish-brown color on the teeth and inside of the mouth. With continued use, these stains become permanent and often come with changes in the mouth lining. Use of smokeless tobacco in various forms like betel

quid, paan, naswar, and areca nut has been prevalent for centuries in South Asian culture [3]. However, Ghutka and Mawa became more popular during the last two decades because of their cheap cost, ready availability in small sachets, and lack of awareness regarding their harmful effects [4]. The method of intake is simply a pinch placed between the cheek and gum, slowly sucked and chewed, very similar to other forms of tobacco chewing. But unlike traditional forms, Ghutka/Mawa is highly addictive and strongly carcinogenic [5]. Several studies have confirmed the association of Ghutka and Mawa with multiple oral and

pharyngeal diseases. These include dental caries, gingival recession, oral ulcers, leukoplakia, oral submucous fibrosis, and malignancies of the oral cavity and pharynx[6-7]. The International Agency for Research on Cancer (IARC) classifies areca nut and smokeless tobacco products as Group 1 carcinogens, with a well-documented role in squamous cell carcinoma of the oral cavity[8]. In Pakistan and India, oral cancers are among the most common malignancies, with strong epidemiological links to Ghutka and Mawa consumption [9-10]. The burden is increasing especially in younger age groups, which makes the problem even more alarming for clinicians working in ENT and oral surgery units.

#### METHODS

This cross-sectional observational study was conducted in the outpatient department of Al-Tibri Medical College and Hospital, Karachi. The study duration was six months, from January 2025 to June 2025. Ethical approval was obtained from the Institutional Review Board (Ref. No. IREC/ATMC/16(02-2024)/01) of Al-Tibri Medical College and Hospital. Written informed consent was taken from every patient before participation. A total of 100 patients were enrolled by non-probability purposive sampling. Sample size was calculated using Cochran's formula for proportions,  $n = \frac{Z^2 \cdot p \cdot q}{e^2}$  where n is the required sample size, Z is 1.96 at 95% confidence level, p is the expected prevalence of oral lesions in Ghutka/Mawa users from earlier studies, q = 1-p, and e (margin of error) was taken as 0.05. The minimum sample size came to around 96, but we included 100 patients to make the results more reliable. Patients above 18 years, with a history of Ghutka/Mawa use for at least six months and presenting with oral complaints, were included. Patients with other forms of tobacco use, systemic illnesses causing oral lesions, or those who refused consent were excluded. Data collection was done by taking a short history mainly regarding Ghutka/Mawa consumption habits like duration, frequency, and any combinations. After that, a detailed oral cavity examination was carried out by the investigator, and findings were noted in a structured Performa. In a few patients, although they came for other medical problems, they themselves talked about the Ghutka habit when examined. Data were entered and analyzed using SPSS version 22.0. Descriptive statistics were applied for demographic and clinical variables. Association between Ghutka/Mawa use and oral cavity findings and duration was checked using the Chisquare test, and a p-value  $\leq 0.05$  was taken as significant. Data collection was done by taking a short history mainly regarding Ghutka/Mawa consumption habits like duration, frequency, and any combinations. The data collection instrument was a self-designed, structured Performa. To ensure content validity, a panel of three senior ENT specialists and a dental public health expert reviewed it. They had their say in order to enhance the simplicity fand understandability of items. To determine the reliability and feasibility of the Performa, 10 patients (excluding the main study) were piloted on it. The pilot demonstrated the steady knowledge and use of the Performa, and it was not necessary to make any significant change. After that, a detailed oral cavity examination was carried out by the investigator, and findings were noted in this structured Performa.

#### RESULTS

Out of 100 patients examined, dental problems were the most common complaint (73 cases), followed by trismus (44), pain (40), and chewing difficulty (21). The majority of patients had addiction for 6–20 years, while 11 patients used it for more than 20 years (Table 1).

**Table 1:** Frequencies of Different Complaints Among Patients and Duration of Addiction

Category	Subcategory	Frequency (n)	
Complaints	Chewing	21	
	Trismus	44	
	Dental Issue	73	
	Pain	40	
Duration of Addition	Less than 5 years	25	
	6-10 years	30	
	11-20 years	34	
	More than 20 years	11	

Ghutka and Mawa showed the highest association with oral lesions. Ulcers were seen in 58 Mawa and 53 Ghutka users, growths in 35 and 29, respectively, while dental issues were frequent in both groups. Other products like niswar, tobacco, pan, and smoking also showed a significant relation, though less frequent. Chi-Square test was applied; level of significance was <0.05 (Table 2).

**Table 2:** Association of Material of Addiction with Examination Findings

Substances	N	Ulcer (N)	Growth (N)	Dental Issue (N)	p-Value
Gutka	53	35	12	25	0.037
Mawa	58	29	22	15	0.016
Pan	8	2	1	5	0.049
Niswar	10	1	0	0	0.001
Tobacco	15	10	2	3	0.001
Smoking	13	2	1	11	0.001
Supari	15	0	0	15	0.001
Manpuri	1	1	0	0	0.001

With longer duration of use, more severe findings were noted. Less than 5 years of habit showed mostly ulcers and dental issues, while patients with more than 10 years of use had a higher frequency of growths, some suspicious of malignancy. The relation between duration and oral

findings was statistically significant. Chi-Square test was applied; level of significance was < 0.05 (Table 3).

**Table 3:** Association of Examination Findings with the Duration of Addiction

Duration of Habit	N	Ulcer (N)	Growth (N)	Dental Issue (N)	p-Value
Less than 5 years	25	2	00	22	0.001
6-10 years	30	11	01	18	0.027
11-20 years	34	13	05	16	0.041
More than 20 years	11	6	4	1	0.001

## DISCUSSION

The patients who took Ghutka and Mawa in our study had oral complaints that ranged from stains in the teeth and mucosa to actual ulcers and fibrotic alterations. These results can be compared to the previous research in Pakistan and India, the long-term smokeless tobacco use has been reported to induce oral submucous fibrosis, leukoplakia, and even squamous cell carcinoma [11-12]. This mechanism is established, and the alkaloids of the areca nut and tobacco are carcinogens; slaked lime increases the penetration of these carcinogens into the mucosa [13]. Among the notable findings was the fact that a good number of patients were not quite aware of the deleterious consequences of Ghutka/Mawa. This lack of awareness and the low cost render it a significant social health concern, particularly among the low socioeconomic populations. These patterns have been recorded in the research of Karachi and other South Asian cities where awareness level regarding the risk of oral cancer is low, and the consumption has been high [14,15]. The results of this paper corroborate the evidence of Ghutka/Mawa being an addictive substance. Admitted patients stated that they could not guit the habit even knowing about the stains and frequent oral ulcers. This addiction is congruent with the pharmacological impact of arecoline (areca nut) and nicotine (tobacco) that are associated with the development of dependence [16]. The addition of paraffin wax and flavoring substances can also be seen as increasing the likability of the product amongst younger users, thus the popularity of adolescents as new consumers can be reported more often [17]. As an ENT, these habits are not restricted to the pathology of the oral cavity. They also diffuse to pharynx and larynx. Past researches have revealed an association with cancers of the oropharynx, trismus secondary to submucous fibrosis and chronic pharyngitis [18-19]. Most of our patients came with irrelevant ENT issues, but when we examined them, we noticed lesions that were clearly a result of Ghutka use. The rising rate of oral cancer cases in Pakistan is also another problem. As Virani et al. and other reports have demonstrated many times over, oral cavity cancers are one of the leading cancers in both men and women, and they are

primarily associated with the use of smokeless tobacco [20]. This is also supported by our findings and indicates that urgent measures to have been taken in the area of public health. It can be reduced through simple measures such as awareness campaigns, preventing sales around schools and ensuring close scrutiny of sachet manufacturing, primarily among the youth. Our study has several strengths, including the fact that it is limited by a small sample, and it was conducted in a single center. Nevertheless, the findings are not in vain as they reflect a clinical trend that ENT doctors have to deal with on a daily basis. The use of Ghutka and Mawa has a close relationship with oral and throat issues. Even when other complaints are presented to the ENT specialists, they must remain alert to observe early mucosal changes to provide early advice and referrals. Moreover, the non-probability sampling in one hospital OPD has some bias in that the study as well may not reflect the entire community of Ghutka and Mawa users as they may be symptom-free, or are not visiting hospitals. Catching of Ghutka and Mawa related lesions at an early stage should be conducted by regularly examining the mouth in ENT and dental clinics. The dangers of these products should be demonstrated through awareness campaigns, mostly among the younger people. This habit can be prevented with the assistance of schools and community programs. The laws which regulate the selling of Ghutka and Mawa need to be tough and a total ban implemented to check this growing health hazard.

## CONCLUSIONS

Our study shows that Ghutka and Mawa use is highly linked with many mouth problems like ulcers, bad teeth, trismus, and some doubtful growths. The severity was more in patients who were using them for long years, mostly above 10 years. Most of the users had very little awareness and visited ENT OPD very late, where we found hidden mouth lesions during a normal checkup. Early finding and good counseling to patients are very important to stop further damage. Public health knowledge and strong control on selling are needed to reduce this bad and growing habit.

## Authors Contribution

Conceptualization: VKK Methodology: VKK, HM Formal analysis: TMG

Writing review and editing: KI, DK, RK

All authors have read and agreed to the published version of the manuscript

## Conflicts of Interest

All the authors declare no conflict of interest.

# Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

## REFERENCES

- [1] Gupta B and Johnson NW. Systematic Review and Meta-Analysis on the Association between Smokeless Tobacco and Betel Quid without Tobacco with the Incidence of Oral Cancer in South Asia and the Pacific. PLoS One. 2014 Nov; 9(11): e113385. doi: 10.1371/journal.pone.0113385.
- [2] Boffetta P, Hecht S, Gray N, Gupta P, Straif K. Smokeless Tobacco and Cancer. The Lancet Oncology. 2008 Jul; 9(7): 667–75. doi: 10.1016/S1470-2045(08)70173-6.
- [3] Khan MA, Vichayanrat T, Ngoenwiwatkul Y. Association between Smoking and Smokeless Tobacco Use with Dental Caries among Pakistani Patients. BMC Oral Health. 2024 Jun; 24(1): 723. doi: 10.1186/s12903-024-04508-y.
- [4] Gajendra S, McIntosh S, Ghosh S. Effects of Tobacco Product Use on Oral Health and the Role of Oral Healthcare Providers in Cessation: A Narrative Review. Tobacco-Induced Diseases. 2023 Jan; 21: 12. doi: 10.18332/tid/157203.
- [5] Senevirathna K, Pradeep R, Jayasinghe YA, Jayawickrama SM, Illeperuma R, Warnakulasuriya S, et al. Carcinogenic Effects of Areca Nut and Its Metabolites: A Review of Experimental Evidence. Clinics and Practice. 2023 Feb; 13(2): 326-46. doi: 10.3390/clinpract13020030.
- [6] Rumgay H, Nethan ST, Shah R, Vignat J, Ayo-Yusuf O, Chaturvedi P, et al. Global Burden of Oral Cancer in 2022 Attributable to Smokeless Tobacco and Areca Nut Consumption: A Population-Attributable Fraction Analysis. The Lancet Oncology. 2024 Nov; 25(11): 1413–23. doi: 10.1016/S1470-2045(24)00458-3.
- [7] Javed F, Chotai M, Mehmood A, Almas K. Oral Mucosal Disorders Associated with Habitual Gutka Usage: A Review. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology. 2010 Jun; 109(6): 857-64.
- [8] Khan MF, Hayhoe RP, Kabir R. Exploring the Risk Factors for Oral Cancer in Pakistan: A Systematic Literature Review. Dentistry Journal. 2024 Jan; 12(2): 25. doi: 10.3390/dj12020025.
- [9] Kazemi K, Fadl A, Sperandio FF, Leask A. The Areca Nut and Oral Submucosal Fibrosis: A Narrative Review. Dentistry Journal. 2025 Aug; 13(8): 364. doi: 10.3390/dj13080364.
- [10] Pimolbutr K, Poomsawat S, Na-ek N, Warnakulasuriya S, Buajeeb W. Prevalence of Human Papillomavirus in Oral Cancer across Asia: A Systematic Review and Meta-Analysis. Oral Diseases. 2025 May; 31(5): 1479-89. doi: 10.1111/odi.14979.
- [11] Pervez S, Khan MR, Ujjan ID, Mumtaz N, Alam E,

- Qureshi H. Hyderabad (Sindh, Pakistan) Cancer Registry Report over Four Years (2020–2023). Small. 2025 May; 45(2.66): C16.
- [12] Niaz K, Maqbool F, Khan F, Bahadar H, Hassan FI, Abdollahi M. Smokeless Tobacco (paan and gutkha) Consumption, Prevalence, and Contribution to Oral Cancer. Epidemiology and Health. 2017 Mar 9; 39: e2017009.
- [13] Dembicka-Mączka D, Fiegler-Rudol J, Skaba D, Kawczyk-Krupka A, Wiench R. High-Energy Lasers in Oral Oncology: A Systematic Review and Meta-Analysis. Journal of Clinical Medicine. 2025 Sep; 14(18): 6419. doi: 10.3390/jcm14186419.
- [14] Sultan Y, Salman Z, Alzaatreh M, Edilbi A, Alani R, Sultan I, et al. Smoking-Related Disease Impact in the Eastern Mediterranean Region: A Comprehensive Assessment Using Global Burden of Disease Data. Asian Pacific Journal of Cancer Prevention. 2024 Feb; 25(2): 495. doi: 10.31557/APJCP.2024.25.2.495.
- [15] Matariya R, Vadera H, Jain S, Vaishnav K, Shah M, Shah R, et al. Influence of Smokeless Tobacco on the Development of Oral Submucous Fibrosis among Industrial Workers: A Cross-Sectional Study. Journal of Datta Meghe Institute of Medical Sciences University. 2023 Jan; 18(1): 39–43. doi: 10.4103/jdmimsu.jdmimsu\_535\_22.
- [16] Anwar N, Chundriger Q, Awan S, Moatter T, Ali TS, Abdul Rasheed M, et al. Prevalence of High-Risk Human Papillomavirus in Oral Squamous Cell Carcinoma with or without Chewing Habits. PLoS One. 2024 May; 19(5): e0300354. doi: 10.1371/journal. pone.0300354.
- [17] Berrone M, Lajolo C, De Corso E, Settimi S, Rupe C, Crosetti E, et al. Cooperation between ENT Surgeons and Dentists in Head and Neck Oncology. Acta Otorhinolaryngologica Italica. 2021 May; 41(2 Suppl 1): S124. doi: 10.14639/0392-100X-suppl.1-41-2021-13.
- [18] von Weymarn LB, Thomson NM, Le Marchand L, Murphy SE. CYP2A6 Activity and Deuterated 4-(Methylnitrosamino)-1-(3-PyridyI)-1-Butanone (NNK) Metabolism in Cigarette Smokers. Chemical Research in Toxicology. 2025 Sep; 38(9): —. doi: 10.1021/acs.chemrestox.5c00310.
- [19] Stoopler ET, Villa A, Bindakhil M, Díaz DL, Sollecito TP. Common Oral Conditions: A Review. Journal of the American Medical Association. 2024 Mar; 331(12): 1045–54. doi: 10.1001/jama.2024.0953.
- [20] Virani SS, Ahmed KS, Springer M, Hussain M, Christensen L, Asif F, et al. Cancer Registries in Pakistan: A Scoping Review. The Lancet Regional Health Southeast Asia. 2025 Jul; 38: 100615. doi: 10.1016/j.lansea.2025.100615.