



Case Study

Prevalence and Predictors of Non-Compliance with Pharmacological Treatment in Diabetes

 Ayesha Shah Jehan¹, Summaya Roshan^{1*}, Shujaat Hussain², Azka Maheen³, Jawad Mahmood⁴ and Akhtar Zada⁵
¹Department of Medicine, Combined Military Hospital, Nowshera, Pakistan

²Department of Gastroenterology, Pakistan Institute of Medical Sciences, Islamabad, Pakistan

³Mufti Mehmood Teaching Hospital, Gomal Medical College, DI Khan, Pakistan

⁴Department of Gastroenterology, Good Hope Hospital, University Hospital, Birmingham, United Kingdom

⁵Department of Medicine, Hayatabad Medical Complex, Peshawar, Pakistan

ARTICLE INFO

Key Words:

Diabetes Mellitus (DM), Non-Adherence, Pharmacological Therapy

How to Cite:

 Jehan, A. S., Roshan, S., Hussain, S., Maheen, A., Mahmood, J., & Zada, A. (2023). Prevalence and Predictors of Non-Compliance with Pharmacological Treatment in Diabetes : Non-Compliance with Pharmacological Treatment in Diabetes. Pakistan Journal of Health Sciences, 4(10). <https://doi.org/10.54393/pjhs.v4i10.1089>

*Corresponding Author:

 Summaya Roshan
 Department of Medicine, Combined Military Hospital,
 Nowshera, Pakistan
Summayaroshan496@gmail.com

 Received Date: 3rd October, 2023

 Acceptance Date: 23rd October, 2023

 Published Date: 31st October, 2023

ABSTRACT

Although the incidence of failure to comply to prescribe drugs is very varied, it may be the most frequent cause of suboptimal treatment results in diabetic patients. In the research region, there are little data on the extent of non-adherence to medicines and related causes. **Objective:** To determine the prevalence of non-adherence to medications in patients with diabetes and the factors leading to such practices. **Methods:** A descriptive study designed was chosen to evaluate 191 patients with diabetes enrolled at the department of medicine, combined military hospital, Nowshera, during the period May 2022 to May 2023. Non-adherence was confirmed clinically by taking history and biochemically by poor glycemic profile. SPSS version 24 was used to analyze the data. **Results:** The age of the patients ranged from 18 to 60 years. The mean age of the patients was 36.17 years with standard deviation 7.53. Age group 18 to 40 years comprised of 113 (59.2%) participants. The male participants were 129 (67.5%) in number. Non-adherence was observed in 86 (45.5%) patients. Factors leading to non-adherence included financial constraints 32 (37.2%), medications related to complications 16 (18.6%). **Conclusions:** Diabetes patients often fail to comply with their prescribed prescriptions in the study region. To reduce disengagement among diabetes patients, public health programs should be reinforced.

INTRODUCTION

Diabetes is a serious health issue with a rising incidence that is characterized by persistent metabolic illness and increased blood glucose levels. Along with rising rates of body fat and cardiovascular illnesses, it is acknowledged by the globe Health Organization (WHO) as being one of the most significant public health issues in the globe for early detection, treatment, and prevention. On a worldwide scale, diabetes is projected to affect 422 million individuals. Citizens of low- and middle-income nations make up the majority of these diabetes patients. Diabetes

is specifically responsible for 1.5 million yearly fatalities [1, 2]. This chronic illness requires a lifetime of therapy, dramatically raising the possibility of severe, long-term effects. It is difficult for hospitals where controlling acute infections is given higher priority, to provide long-term surveillance and treatment in cases of diabetes [3]. Diabetes mellitus is an unsettling health issue, despite the fact that its incidence in the area is currently lower than that of affluent nations [4, 5]. According to the most current meta-analysis research the prevalence of diabetes

mellitus in Ethiopia is thought to be 6.5%, with regional differences in the percentage [6, 7]. Medication non-compliance encompasses a broad variety of behaviours, whether they are unintentionally engaged in or not, that either result in the inadequate utilization or misuse of prescription drugs [8]. Breaking tablets, missing dosages, discontinuing medicine early, and failing to renew a prescription are all indications of insufficient utilization of pharmaceuticals. Non-adherence to anti-diabetic treatment is often linked to poor metabolic control. Due to inadequate metabolic regulation, both immediate and long-term problems may arise [9]. Studies have shown that diabetic individuals had suboptimal compliance to their anti-diabetic drugs and hyperglycemia control [10]. The most frequent cause of deteriorating health among diabetics may be failure to comply with treatment. Poor patient adherence compromises the safety and efficacy of anti-diabetic drugs, increasing the risk of death and disability. This has an immediate or subsequent effect on the rising expenses of the healthcare system [11]. Anti-diabetic drug failure to comply percentages are known to be very varied; as a result, the effects on outcomes of therapy should be taken into account [12]. Many diabetes individuals take fewer of their prescriptions than are recommended for them, it has been observed. Thus, increasing medication adherence for diabetes is a critical public health concern [13]. One of the most often mentioned factors influencing medication non-adherence to diabetes therapy is age, followed by medication awareness and the presence of co-morbidities [14]. Designing more effective treatments requires knowing how often diabetes patients don't take their prescriptions as prescribed and the accompanying causes. The prevalence of medication non-adherence, related variables, and its effects on treatment outcomes among diabetes patients in the research region are not described in any studies. Consequently, the purpose of this research was to evaluate medication non-compliance and related factors among diabetes patients.

METHODS

This descriptive analysis was carried out Combined Military Hospital, Nowshera, over the period of May 2022 to May 2023. The hospital is situated in the town of District Nowshera, located in the Khyber Pakhtunkhwa Province of Pakistan. The organization offers services throughout four primary clinical departments, namely internal medicine, surgery, pediatrics, and gynecology/obstetrics. Additionally, it provides a range of services in additional clinical departments including dentistry, orthopedics, diabetes, dermatology, and reproductive health. The study population comprised of patients visiting diabetes clinic with poor glycemic profile. Poor glycemic profile was

defined by persistently raised random blood glucose level >200mg/dl for two weeks. Both type 1 and type 2 diabetes patients in the age range 18 to 60 years were registered. Participants of either gender were enrolled. Non-compliance was defined by failure on the part of the patient to take antidiabetic medications in the prescribed frequency or strength deliberately [15]. Multi-morbidity including intellectual disabilities, failure to provide formal authorization, serious auditory and vision impairments, lack of willingness and patients with severe cardiopulmonary compromise were excluded [16]. Non probability consecutive sampling technique was employed. A total of 191 patients were recruited. Sample size was determined using WHO sample size formula taking the anticipated prevalence of non-compliance as 41.5%, margin of error 7% and confidence level 95% [15]. In order to make sure that patient rights were upheld and ethical norms were met, the research plan was presented to the research ethics committee of the hospital. Prior to starting data collection, ethical approval was obtained, and patient data were concealed to maintain confidentiality. Data were collected from outdoor diabetes clinic of the hospital. Patients with poor glycemic profile were registered. Detailed history was taken from the patient regarding the frequency and the strength in which medications were being taken. These observations were then compared with the prescribed frequency and strength. Deliberate reduction in the frequency or strength of the medications was noted. History was further elaborated to discover the factors leading to poor adherence. Software version 25.0 of the Statistical Package for Social Sciences (SPSS) was used to code and provide data for analysis. To provide an overview of categorical variables, descriptive statistics like frequencies and percentages were calculated and the results were shown as tables. The relationship between the predictors and the result variable was examined using binary logistic regression models. Statistical significance was defined as p -value ≤ 0.05 at 95% CI.

RESULTS

The age of the patients ranged from 18 to 60 years. The mean age of the patients was 36.17 years with standard deviation of 7.53. Frequency of participants in the age group 18 to 40 years was 113 (59.2%) while rest of the patients 78 (40.8%) had age between 41 and 60 years. The number of male participants in the study was 129 (67.5%). Residence wise distribution of patients showed that 119 (62.3%) patients belonged to rural areas while the remaining 72 participants (37.7%) were living in urban areas. The monthly income of 140 patients (73.2%) was more than Rs. 50,000 per month. The mean duration of disease was 8.57 years with standard deviation 3.91. Majority of the participants (110, 57.6%) had disease

duration more than 5 years and the remaining 81 patients (42.4%) had disease duration less than 5 years. 175 patients (91.6%) had Type 2 DM while the rest of 16 patients (8.4%) had type 1 diabetes mellitus. 137 patients (71.7%) patients were advised oral antidiabetic medications on their prescription while rest of the 54 patients (28.3%) were advised injectable medication (insulin). Figure 1 illustrates that non adherence was observed in 86 patients (45.0%). Age wise distribution showed 56 patients (49.5%) had age 18 to 40 years while 30 patients (38.5%) were aging 41 to 60 years. Contingency table analysis for non-adherence versus age reveal chi square p value of 0.129 which was more than 0.05. Hence it was statistically not significant. Similarly gender wise non-adherence break up was 69 (53.5%) among male as compared to 16 participants (25.8%) among female participants. The chi square p value was 0.001 which was less than 0.05. Hence, the heterogeneity was statistically significant. 42 patients (77.8%) among those taking injectable oral antidiabetic were non-compliant as compared to 44 patients (32.1%) taking oral hypoglycemic were non-compliant. The chi square p value was 0.001 which was less than 0.05, hence it was statistically significant.




Figure 1: Distribution of patients with respect to non-compliance Table 1 shows that financial constraints were reported by 32 participants (37.2%), followed by fear of injection/injection related apprehensions in 20 patients (23.2%). Other causes included bad experience with medications such as adverse effects reported by 16 patients (18.6%), misconceptions about treatment described by 07 patients (8.1%) and lack of knowledge was observed in 11 participants (12.8%) who were non-complaint to medications.

Table 1: Factors leading to non-compliance

Factors	Frequency (%)
Financial Constraints	32 (37.2)
Injection Fear/Apprehensions	20 (23.3)
Medications adverse effects	16 (18.6)
Misconceptions	07 (8.1)
Lack of Knowledge	11 (12.8)

DISCUSSION

We discovered that 86 (45.5%) of the respondents in our study were non-adherent. The percentage of non-adherence to treatment in our research was comparable to the results of studies. Mitiku *et al.*, reported 41.0% non-adherence rate in their study [15]. Ali *et al.*, reported 45.2% non-compliance to antidiabetic medications [17]. Nevertheless, the observations in this research were substantially greater compared with the findings of research projects carried out by Bonger *et al.*, [18]. Nandini *et al.*, reported 21.6% non-adherence rate in a study carried out on Indian population [19]. In contrast, the results for non-adherence to antidiabetic medications were higher than the observations of this study [20]. This variability in the results of different studies may be attributed to the different sample sizes and adherence testing methods that may affect non-adherence levels and contribute to the variations in study outcomes [21]. Additionally, the amount of non-adherence as shown in various research settings may vary according to the participants' educational status, place of residence, income, and degree of medicine knowledge [22]. The majority of patients who sought medical attention in the current study were younger. Age and non-adherence do not significantly correlate, according to the current research (p-value = 0.129). However, non-compliance to medications was more common in patients aging more than 40 years. In a study by Mitiku *et al.*, the rate of non-adherence to medication was twice as common in patients aging more than 40 years compared to patients younger than 40 years [15]. This may be attributed to the fact that forgetfulness increases with age which was shown in a research, according to which the knowledge score drops by 3% for every 10 years increase in age after 40 years [23]. In our study, almost half of the male patients were non-compliant to medication as compared to one fourth females. This difference was statistically significant (p < 0.05). This finding is similar to the result of another study [24]. Imtiaz *et al.*, reported showed no evidence of a link between gender and non-adherence in diabetics [25]. The high rate of non-adherence in men may be because men often participate in outdoor pursuits in our nation, which may lead them to forget to take their prescriptions. Compared to urban inhabitants, rural residents less likely to adhere. This finding is comparable to the finding of Tiv *et al.*, [26]. This may be attributed to the fact that in addition to communication deficiencies, patients in remote locations likely experience other difficulties like convenient access to healthcare services.

CONCLUSIONS

There is a statistically significant high rate of medication non-adherence among DM patients. The higher risk of antidiabetic medication non-adherence was substantially

correlated with male sex, living in a rural area, weak monthly income and patients who were advised injectable medications. Implementing strategies such as mental and emotional support, appropriate counselling and enhanced communication with healthcare experts, assistance from family members, and resolving financial restraints to meet their medical bills are all necessary to raise the adherence rate of diabetes patients. Furthermore, multicenter prospective trials that reduce recollection bias are advised in order to identify the true determinants of poor adherence.

Authors Contribution

Conceptualization: SR, SH, JM

Methodology: ASJ, SR

Formal analysis: ASJ, SR, AM, AZ

Writing-review and editing: SH, JM, AZ

All authors have read and agreed to the published version of the manuscript.

Conflicts of Interest

The authors declare no conflict of interest.

Source of Funding

The authors received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Mann DM, Ponieman D, Leventhal H, Halm EA. Predictors of adherence to diabetes medications: the role of disease and medication beliefs. *Journal of Behavioral Medicine*. 2009 Jun; 32: 278-84. doi: 10.1007/s10865-009-9202-y.
- [2] Nuche-Berenguer B and Kupfer LE. Readiness of sub-Saharan Africa healthcare systems for the new pandemic, diabetes: a systematic review. *Journal of Diabetes Research*. 2018 Feb; 2018: 9262395. doi: 10.1155/2018/9262395.
- [3] Jufar AH, Jima Z, Gameda D. Prevalence and factors contributing to non-adherence to diabetes treatment among diabetic patients attending government hospitals in Addis Ababa. *International Journal of Tropical Disease & Health*. 2018 Oct; 32(3): 1-1. doi: 10.9734/IJTDH/2018/43867.
- [4] Lee WC, Balu S, Cobden D, Joshi AV, Pashos CL. Medication adherence and the associated health-economic impact among patients with type 2 diabetes mellitus converting to insulin pen therapy: an analysis of third-party managed care claims data. *Clinical Therapeutics*. 2006 Oct; 28(10): 1712-25. doi: 10.1016/j.clinthera.2006.10.004.
- [5] Zeru MA, Tesfa E, Mitiku AA, Seyoum A, Bokoro TA. Prevalence and risk factors of type-2 diabetes mellitus in Ethiopia: systematic review and meta-analysis. *Scientific Reports*. 2021 Nov; 11(1): 21733. doi: 10.1038/s41598-021-01256-9.
- [6] World Health Organization. Adherence to long-term therapies: evidence for action. World Health Organization; 2003.
- [7] Adisa R, Alutundu MB, Fakeye TO. Factors contributing to nonadherence to oral hypoglycemic medications among ambulatory type 2 diabetes patients in Southwestern Nigeria. *Pharmacy Practice*. 2009 Jul; 7(3): 163. doi: 10.4321/S1886-36552009000300006.
- [8] García-Pérez LE, Álvarez M, Dilla T, Gil-Guillén V, Orozco-Beltrán D. Adherence to therapies in patients with type 2 diabetes. *Diabetes Therapy*. 2013 Dec; 4: 175-94. doi: 10.1007/s13300-013-0034-y.
- [9] Kasznicki J, Głowacka A, Drzewoski J. Type 2 diabetic patients compliance with drug therapy and glycaemic control. *Diabetologia Doświadczalna i Kliniczna*. 2007 Jul; 7(4): 199-203.
- [10] Boshe BD, Yimar GN, Dadhi AE, Bededa WK. The magnitude of non-adherence and contributing factors among adult outpatient with Diabetes Mellitus in Dilla University Referral Hospital, Gedio, Ethiopia. *PloS One*. 2021 Mar; 16(3): e0247952. doi: 10.1371/journal.pone.0247952.
- [11] Abate TW. Medication non-adherence and associated factors among diabetes patients in Felege Hiwot Referral Hospital, Bahir Dar city administration, Northwest Ethiopia. *BMC Research Notes*. 2019 Dec; 12(1): 1-6. doi: 10.1186/s13104-019-4205-4.
- [12] Demoz GT, Berha AB, Alebachew Woldu M, Yifter H, Shibeshi W, Engidawork E. Drug therapy problems, medication adherence and treatment satisfaction among diabetic patients on follow-up care at Tikur Anbessa Specialized Hospital, Addis Ababa, Ethiopia. *PloS One*. 2019 Oct; 14(10): e0222985. doi: 10.1371/journal.pone.0222985.
- [13] Kusa W, Tolessa D, Abdeta T. Type II DM medication non-adherence in Adama Hospital Medical College, Central Ethiopia. *East African Journal of Health and Biomedical Sciences*. 2019 May; 3(1): 31-8.
- [14] Kirkman MS, Rowan-Martin MT, Levin R, Fonseca VA, Schmittiel JA, Herman WH, et al. Determinants of adherence to diabetes medications: findings from a large pharmacy claims database. *Diabetes Care*. 2015 Apr; 38(4): 604-9. doi: 10.2337/dc14-2098.
- [15] Mitiku Y, Belayneh A, Tegegne BA, Kebede B, Abebe D, Biyazin Y, et al. Prevalence of Medication Non-Adherence and Associated Factors among Diabetic Patients in A Tertiary Hospital at Debre Markos, Northwest Ethiopia. *Ethiopian Journal of Health*

- Sciences. 2022 Jul; 32(4): 755-764. doi: 10.4314/ejhs.v32i4.12.
- [16] Foley L, Larkin J, Lombard-Vance R, Murphy AW, Hynes L, Galvin E, et al. Prevalence and predictors of medication non-adherence among people living with multimorbidity: a systematic review and meta-analysis. *BMJ Open*. 2021 Sep; 11(9): e044987. doi: 10.1136/bmjopen-2020-044987.
- [17] Ali M, Alemu T, Sada O. Medication adherence and its associated factors among diabetic patients at Zewditu Memorial Hospital, Addis Ababa, Ethiopia. *BMC Research Notes*. 2017 Dec; 10(1): 1-5. doi: 10.1186/s13104-017-3025-7.
- [18] Bongor Z, Shiferaw S, Tariku EZ. Adherence to diabetic self-care practices and its associated factors among patients with type 2 diabetes in Addis Ababa, Ethiopia. *Patient Preference Adherence*. 2018 Jun; 12: 963-970. doi: 10.2147/PPA.S156043.
- [19] Nandini HC, Gali A, Muraraiah S. Assessment of factors influencing adherence to antidiabetic drugs among patients with type 2 diabetes mellitus at a tertiary care hospital in India. *Pharmacology and Clinical Pharmacy Research*. 2020 Apr; 5(1): 7-13. doi: 10.15416/pcpr.v5i1.26507.
- [20] Aminde LN, Tindong M, Ngwasiri CA, Aminde JA, Njim T, Fondong AA, et al. Adherence to antidiabetic medication and factors associated with non-adherence among patients with type-2 diabetes mellitus in two regional hospitals in Cameroon. *BMC Endocrinology Disorders*. 2019 Apr; 19(1): 35. doi: 10.1186/s12902-019-0360-9.
- [21] Araya EM, Gebrezgabihier HA, Tekulu GH, Alema NM, Getnet D, Gebre HT, et al. Medication non-adherence and associated factors among diabetic patients visiting general hospitals in the eastern zone of Tigray, Northern Ethiopia. *Patient Preference and Adherence*. 2020 Oct; 14: 2071-83. doi: 10.2147/PPA.S278148.
- [22] Bagonza J, Rutebemberwa E, Bazeyo W. Adherence to anti diabetic medication among patients with diabetes in eastern Uganda; a cross sectional study. *BMC Health Services Research*. 2015 Dec; 15(1): 1-7. doi: 10.1186/s12913-015-0820-5.
- [23] Shams N, Amjad S, Ahmed W, Saleem F. Drug non-adherence in type 2 diabetes mellitus; predictors and associations. *Journal of Ayub Medical College Abbottabad*. 2016 Jun; 28(2): 302-7.
- [24] Parimalakrishnan S, Dussa K, Sahay R. Assessment of diabetes knowledge using diabetes knowledge questionnaire among people with type 2 diabetes mellitus. *Asian Journal of Pharmaceutical and Clinical Research*. 2015 Mar; 8(2): 254-56.
- [25] Imtiaz S, Ullah H, Rasool MF, Hashmat F, Saleem M, Khan N. Assessment of Compliance of Diabetic Patients at Nishtar Hospital Multan, Pakistan. *Gomal Journal of Medical Sciences*. 2014 Jun; 12(2): 84-8.
- [26] Tiv M, Viel JF, Mauny F, Eschwege E, Weill A, Fournier C, et al. Medication adherence in type 2 diabetes: the ENTRED study 2007, a French population-based study. *PLoS One*. 2012 Mar; 7(3): e32412. doi: 10.1371/journal.pone.0032412.