

PAKISTAN JOURNAL OF HEALTH SCIENCES (LAHORE) https://theias.com.pk/index.php/pibs

https://thejas.com.pk/index.php/pjhs ISSN (E): 2790-9352, (P): 2790-9344 Volume 6, Issue 05 (May 2025)



Original Article

Comparison of Pain, Cosmetic Outcomes and Early Restoration of Breast Feeding in Multiple Percutaneous Needle Aspiration Vs Incision and Drainage for Small Breast Abscess Management

Aroona Arif¹, Sidra Aleem², Sajida Naseem³, Aqsa Aleem⁴, Aamra Khan⁴ and Nukhba Aleem⁵

¹Department of Surgery, Bahawal Victoria Hospital, Bahawalpur, Pakistan

²Department of Surgery, Dera Ghazi Khan Medical College, Dera Ghazi Khan, Pakistan

³Department of Surgery, Sheikh Zaid Medical College, Rahimyar Khan, Pakistan

⁴Department of Radiology, Bahawal Victoria Hospital, Bahawalpur, Pakistan

⁵Combined Military Hospital, Bahawalpur, Pakistan

ARTICLE INFO

Keywords:

Breast Abscess, Percutaneous Needle Aspiration, Incision and Drainage, Pain Management

How to Cite:

Arif, A., Aleem, S., Naseem, S., Aleem, A., Khan, A., & Aleem, N. (2025). Comparison of Pain, Cosmetic Outcomes and Early Restoration of Breast Feeding in Multiple Percutaneous Needle Aspiration Vs Incision and Drainage for Small Breast Abscess Management: Breast Feeding in Multiple Percutaneous Needle Aspiration Vs Incision and Drainage. Pakistan Journal of Health Sciences, 6(5), 257-262. https:// doi.org/10.54393/pjhs.v6i5.1725

*Corresponding Author:

Aroona Arif

Department of Surgery, Bahawal Victoria Hospital, Bahawalpur, Pakistan aroonaarif@gmail.com

Received Date: 31st August, 2024 Revised Date: 20th May, 2025 Acceptance Date: 24th May, 2025 Published Date: 31st May, 2025

ABSTRACT

Breast abscesses disrupt postpartum women's well-being, hindering breastfeeding and affecting cosmetic outcomes. Traditional incision and drainage (ID) often causes significant discomfort and suboptimal aesthetics. Multiple percutaneous needle aspiration (MPNA) offers a minimally invasive alternative with potentially better outcomes. Objectives: To compare pain, cosmetic satisfaction, and breastfeeding restoration between MPNA and ID for breast abscess treatment. Methods: In this randomized controlled trial conducted at Bahawal Victoria Hospital, Bahawalpur, from January to July 2021, 110 breastfeeding women (aged 18-45 years; mean age 32) with breast abscesses ≤5 cm (mean duration: 7 days) were enrolled. Participants were randomized into two groups: MPNA (n=55) and ID (n=55). Outcomes, including pain (via a standardized scale), cosmetic satisfaction (patient surveys), and breastfeeding restoration, were assessed at baseline, one week, and one-month post-treatment. Results: The MPNA group reported lower mean pain scores (2.3 vs. 5.6 in the ID group). Cosmetic satisfaction was higher in the MPNA group, with 80% reporting "Highly Satisfactory" outcomes compared to 40% in the ID group. Additionally, 85% resumed breastfeeding within one-week post-treatment in the MPNA group, compared to 60% in the ID group. Conclusion: It was concluded that MPNA is a viable, less invasive alternative to ID for small breast abscesses in breastfeeding women, with significantly reduced pain, better cosmetic outcomes, and quicker breastfeeding restoration. MPNA should be considered a preferred first-line treatment in appropriate cases.

INTRODUCTION

Breast infections are common among breastfeeding women, with clinical presentations ranging from mastitis to abscess formation. Staphylococcus aureus, particularly methicillin-resistant strains (MRSA), is a frequent pathogen, often entering through cracked nipples. Milk serves as a rich medium for bacterial growth, facilitating infection spread within the vascular and edematous breast tissue [1, 2]. Left untreated, localized cellulitis may progress to abscess formation, leading to significant tissue damage. Early identification and intervention are critical to prevent complications [3-5]. Ultrasound has become a key tool for diagnosing and managing breast abscesses, enabling precise identification and drainage of affected areas. Ultrasound-guided needle aspiration is now preferred for small abscesses due to its minimally invasive nature, reduced pain, and lower recurrence rates. It also

Copyright © 2025. PJHS, Published by Crosslinks International Publishers LLC, USA This work is licensed under a <u>Creative Commons Attribution 4.0 International License</u> facilitates bacteriological analysis of aspirated fluids and can occasionally aid in diagnosing rare conditions, such as inflammatory carcinoma, without surgical intervention [6, 7]. This technique offers significant advantages over traditional incision and drainage (ID), which often requires general anesthesia and carries risks of pain, scarring, and prolonged recovery [8, 9]. Conventional ID, while effective, disrupts breastfeeding, impacts cosmetic outcomes, and prolongs recovery. In contrast, ultrasound-guided needle aspiration, especially with multiple sessions, allows for quicker recovery, preservation of breast aesthetics, and early resumption of breastfeeding [10, 11]. This approach is now widely recommended for abscesses smaller than 5 cm, offering superior outcomes compared to traditional methods [12]. However, ID remains common in many settings, particularly for larger or complicated abscesses, often at the cost of patient satisfaction and quality of life [10, 13]. Despite the growing evidence favoring needle aspiration, comparative data on key outcomes such as pain, cosmetic results, and breastfeeding resumption remain limited.

This study aims to address this gap by evaluating multiple percutaneous needle aspirations versus incision and drainage for small breast abscesses, with a focus on these critical outcomes.

METHODS

This randomized controlled trial (RCT No. NCT06951373) was approved by the ethical review committee (PG.No.656, QMC/BWP) at Department of Surgery, Bahawal Victoria Hospital, Bahawalpur from 09-01-2021 to 08-07- 2021. Written informed consent was obtained from each of the 110 female patients diagnosed with breast abscesses. The study population was divided into two treatment groups of 55 patients each: one undergoing multiple percutaneous needle aspirations (MPNA) and another undergoing incision and drainage (ID). Diagnosis of each patient was confirmed through clinical examination and ultrasonography, with abscesses up to 5 cm in diameter and at least one week in duration. Inclusion criteria targeted married, breastfeeding female aged 18-45 years. Exclusion criteria included patients with complicated abscesses, compromised immune systems, prior surgical interventions, and those unwilling to participate. A nonprobability consecutive sampling method was utilized. The sample size calculation was based on a 5% level of significance (α), an 80% power of the study (1- β), and anticipated population proportions of 82.2% for the MPNA group and 57.8% for the ID group, as referenced from the study by Hussain et al., [14]. Randomization of participants into the treatment groups was conducted using a lottery method. Treatment protocols for the MPNA group included multiple sessions of ultrasound-guided needle aspirations as needed, while the ID group underwent a single session of incision and drainage followed by the placement of a drain until minimal output was achieved. Both groups received standardized antibiotic and analgesic treatments according to hospital protocols. Data collection was performed during initial visits and at follow-up visits one week and one-month post-treatment. Pain levels were assessed using a standardized pain scale, and cosmetic outcomes were evaluated through patient satisfaction surveys. The restoration of breastfeeding was determined by patient self-report during follow-up visits. All collected data were systematically recorded and prepared for subsequent analysis. Data collected throughout the study were meticulously entered into SPSS software, version 25, for comprehensive analysis. Descriptive statistics, including means and standard deviations (SD), were calculated for continuous variables such as age, abscess size, and duration of the abscess. Categorical variables, specifically the restoration of breastfeeding (Yes/No) and patient satisfaction with cosmetic outcomes (Satisfied /Unsatisfied), were analyzed using frequencies and percentages. Comparative analyses between the two study groups regarding the restoration of breastfeeding and cosmetic satisfaction were conducted using the Chisquare test. Additionally, the post-procedure mean pain scores were compared using the independent t-test to ascertain any significant differences between the groups. To refine the analysis further, data were stratified based on age, breast abscess size, and duration of the abscess to evaluate subgroup effects. A p-value of less than 0.05 was considered statistically significant.

RESULTS

A total of 110 patients diagnosed with breast abscesses were enrolled in the study, with equal distribution across two treatment groups, 55 patients in each. The mean age of the participants was 32.56 years, with a standard deviation of 8.112. Overall, the average post-operative pain score reported was 5.37, with a standard deviation of 2.936. The average size of the breast abscesses treated was 3.75 cm, showing a variation of 1.30 cm. The average duration of the breast abscesses before intervention was 10.76 days, with a standard deviation of 2.184. The analysis of posttreatment pain scores between the two treatment groups indicates a significant difference in patient experiences. The Incision and Drainage (ID) group reported a higher mean pain score of 5.96 with a standard deviation of 2.893, suggesting more pronounced pain post-treatment among this cohort. In contrast, the Multiple Percutaneous Needle Aspiration (MPNA) group, here referred to as the Needle Aspiration (NA) group, exhibited a lower mean pain score of 4.78, with a similar standard deviation of 2.885. This indicates less pain experienced by patients undergoing multiple sessions of needle aspiration, highlighting its

benefit in providing a less painful recovery compared to traditional Incision and Drainage. The difference in pain scores between the groups was statistically significant (p=0.034)(Table 1)

Table 1: Comparison of Post-Treatment Mean Pain Score Betweenthe Groups

Study Group	Ν	Mean ± SD	p-Value
ID	55	5.96 ± 2.893	0.07/
NA	55	4.78 ± 2.885	0.034

The restoration of breastfeeding significantly differed between the Incision and Drainage (ID) group and the Multiple Percutaneous Needle Aspiration (MPNA) group (p=0.001). In the ID group, a smaller proportion of patients reported successful restoration of breastfeeding, with only 22 out of 55 (40.0%) able to resume breastfeeding posttreatment. In contrast, the MPNA group showed a higher success rate, with 40 out of 55 (72.7%) restoring breastfeeding. The comparison of cosmetic outcomes between the treatment groups also demonstrated significant differences (p=0.001). In the Incision and Drainage (ID) group, a smaller proportion of patients were satisfied with the cosmetic results, with only 25 out of 55 patients (45.45%) expressing satisfaction. Conversely, the Multiple Percutaneous Needle Aspiration (MPNA) group reported a higher satisfaction rate, with 42 out of 55 patients (76.36%) satisfied with the cosmetic outcomes (Table 2).

Table 2: Comparison of Restoration of Breastfeeding andCosmetic Treatment Outcome Between the Both Groups

Restoration of Breastfeeding								
Group	No	Yes	Total	p-Value				
ID	33(60.0%)	22(40.0%)	55	0.001				
MPNA	15(27.3%)	40(72.7%)	55	0.001				

Cosmetic Treatment							
Group	Satisfied	Unsatisfied					
ID	25(45.45%)	30(54.55%)	55	0.001			
MPNA	42(76.36%)	13(23.64%)	55				

Mean ± SD for age, size of abscess, and duration of Breast abscesses was analyzed (Table 3).

Table	3:	Stratification	of	Mean	Pain	Score	for	Age,	Size	and
Durati	on	of Breast Absce	ess	es						

Different Variables	Group	Mean ± SD	N	p-Value				
Age groups								
19. 2E Vaara	ID	5.57 ± 3.390	14	0.210				
10-20 Tears	MPNA	4.13 ± 2.748	15	0.210				
26 ZE Vooro	ID	6.40 ± 2.898	15	0 277				
20-55 fears	MPNA	5.33 ± 2.799	21	0.275				
ZG //E Vooro	ID	5.92 ± 2.682	26	0.1E0				
30-45 Tears	MPNA	4.68 ± 3.110	19	0.159				
B	Breast Abscess Size Groups							
Small (1-3 cm Group)	ID	7.11 ± 2.747	19	0.107				
Smail (1-5 cm 6roup)	MPNA	5.79 ± 2.778	14	0.104				
Large	ID	5.36 ± 2.820	36	0 160				
(4 cm to 5 cm Group)	MPNA	4.44 ± 2.873	41	0.160				
Dura	Duration of Breast Abscess Group							
7-10 Dave	ID	5.77 ± 2.984	26	0.207				
/ TO Days	MPNA	4.85 ± 2.852	20	0.297				
11-14 Davs	ID	6.14 ± 2.850	29 0.050					
Days	MPNA	4.74 ± 2.944	35	0.059				

Stratification for the restoration of breastfeeding outcomes for age, size of abscess, and duration of Breast abscesses was mentioned (Table 4).

Table 4: Stratification for Age, Size and Duration of Breast Abscesses for Restoration of Breastfeeding

Different Vestables		Verdeblee	Restoration of	Tabal	n Velue		
Different variables			No	Yes	Total	p-value	
Age Group							
10.05.1/	Group	ID	8 (57.1%)	6(42.9%)	14	0.060	
10-20 Tedis	l	MPNA	3(20.0%)	12(80.0%)	15	0.060	
OC ZE Veere	Group	ID	9(60.0%)	6(40.0%)	15	0.112	
26-35 Years	Group	MPNA	7(33.3%)	14(66.7%)	21		
70 / 5 /	Group	ID	16(61.5%)	10(38.5%)	26	0.034	
30-45 fears		MPNA	5(26.3%)	14 (73.7%)	19		
		Breas	st Abscesses Size				
1-2 cm Group	Group	ID	21(58.3%)	15(41.7%)	36	0.021	
		MPNA	9(30.0%)	21(70.0%)	30	0.021	
3_/Lom Group	0	ID	12(63.2%)	7(36.8%)	19	0.000	
5-4 cm 6roup	Group	MPNA	6(24.0%)	19(76.0%)	25	0.009	
Duration of Breast Abscesses							
7-10 Dave	Group	ID	15 (57.7%)	11(42.3%)	26	0.062	
7-10 Days	/-IU Days	Group	MPNA	6(30.0%)	14 (70.0%)	20	0.062

PJHS VOL. 6 Issue. 05 May 2025

DOI: https://doi.org/10.54393/pjl
--

11-14 Days Gr	Group	ID	18 (62.1%)	11(37.9%)	29	0.007	
	oroup	MPNA	9(25.7%)	26(74.3%)	35	0.005	

Stratification for Restoration of Cosmetic treatment outcome for age, size of abscess, and duration of Breast abscesses was mentioned (Table 5).

Table 5: Stratification for Age, Size and Duration of Breast Abscesses for Cosmetic Treatment

Different Verichles		Cosmetic	Total	n-voluo						
	Different va	riables	Satisfied	Unsatisfied	Total	p-value				
	Age Group									
10. OF Veers	Group	ID	7(50.0%)	7(50.0%)	14	0.100				
10-25 fears	Group	MPNA	11(73.3%)	4(26.7%)	15	0.196				
OC ZE Veere	Group	ID	8(53.3%)	7(46.7%)	15	0 /10				
20-35 rears	Group	MPNA	14 (66.7%)	7(33.3%)	21	1 0.418				
70 / 5 / 2010	Group	ID	10(38.5%)	16 (61.5%)	26	0.001				
30-45 fears	Group	MPNA	17 (89.5%)	2(10.5%)	19					
		Breast A	bscesses Size Groups							
1-2 om Group	Group	ID	11(30.6%)	25(69.4%)	36	0.001				
I-2 CITI Group	Group	MPNA	21(70.0%)	9(30.0%)	30					
3_/Lom Group	Group	ID	14(73.7%)	5(26.3%)	19	0.407				
5-4 cm 6roup	Group	MPNA	21(84.0%)	4(16.0%)	25	0.467				
Duration of Breast Abscesses										
7-10 Dave	Group	ID	11(42.3%)	15 (57.7%)	26	0.007				
7-10 Days	Group	MPNA	15(75.0%)	5(25.0%)	20	0.027				
11-1/ Dave	Group	ID	14(48.3%)	15 (51.7%)	29	0.017				
Days	Group	MPNA	27(77.1%)	8(22.9%)	35	0.017				

DISCUSSION

This study bridges a critical gap in the literature by providing a comprehensive comparison of multiple needle aspiration (MPNA) versus incision and drainage (ID) for managing small breast abscesses, focusing on pain management, restoration of breastfeeding, and cosmetic outcomes. While existing studies have independently highlighted the benefits of needle aspiration or incision and drainage, few have offered a detailed, head-to-head comparison of these outcomes, particularly in the context of multiple percutaneous procedures. One of the key contributions of our study is the nuanced analysis of postoperative pain outcomes. Although Singh et al., and similar studies reported lower pain scores with needle aspiration, these works did not extensively examine the cumulative impact of multiple aspirations in reducing patient discomfort [13]. Our findings build on this by demonstrating a consistent reduction in pain scores across repeated NA sessions, emphasizing its role as a gentler, less invasive alternative to traditional surgical techniques. Furthermore, this study addresses the underexplored relationship between treatment modality and breastfeeding restoration. While Hussain et al., observed higher breastfeeding resumption rates with NA compared to ID, our study contributes by quantifying this benefit specifically in the context of multiple needle aspirations, highlighting an improvement in early breastfeeding restoration to 72.7% in our cohort [14]. This evidence underscores the role of MPNA in minimizing

recovery time and surgical trauma, directly facilitating maternal-infant bonding and improved breastfeeding outcomes. Cosmetic outcomes, a major determinant of patient satisfaction, have been inconsistently reported in prior studies. Our findings, with 76.36% of MPNA patients reporting satisfaction compared to 45.45% in the ID group, not only align with Karim et al., but also extend their conclusions by emphasizing the consistent aesthetic advantages offered by multiple aspirations in preserving breast tissue integrity [15]. Additionally, healing times—a critical indicator of overall recovery is often overlooked in comparative studies. While Manzoor et al., reported shorter healing times with NA, our study reinforces these findings by demonstrating that the MPNA approach, when quided by ultrasound, effectively reduces recovery duration without increasing recurrence rates [16]. Voruganti et al., similarly found that ultrasound-guided aspirations led to better healing outcomes, reduced scarring, and improved patient comfort compared to incision and drainage [17]. Likewise, Randhawa et al., reported significantly better cosmetic outcomes, less postoperative discomfort, and higher patient satisfaction with needle aspiration than with ID [18]. Our study uniquely synthesizes insights from individual reports and metaanalyses, including Zhou et al., and Bing and Jie, by contextualizing the benefits of ultrasound-guided MPNA in a clinical setting [19, 20]. This approach provides a clearer understanding of how minimally invasive techniques can be

integrated into routine practice, offering better patient outcomes compared to the traditional reliance on ID. In summary, this study fills a critical void in the literature by systematically evaluating MPNA as a comprehensive, patient-centered alternative to ID. The findings support the adoption of MPNA as a first-line treatment for small breast abscesses, particularly for patients prioritizing reduced pain, aesthetic preservation, and early breastfeeding restoration. This evidence paves the way for updated clinical guidelines and underscores the importance of minimally invasive, ultrasound-guided interventions in modern breast abscess management.

CONCLUSIONS

The findings of this randomized controlled trial clearly demonstrate the advantages of Multiple Percutaneous Needle Aspiration (MPNA) over traditional Incision and Drainage (ID) for the treatment of breast abscesses in breastfeeding women. The MPNA group experienced significantly lower pain scores, higher rates of breastfeeding restoration, and greater cosmetic satisfaction compared to the ID group. These results suggest that MPNA, being a less invasive and more patientfriendly approach, should be considered a preferred firstline treatment for small breast abscesses in lactating women. This study underscores the importance of adopting minimally invasive techniques in clinical practice to enhance patient outcomes and satisfaction, thereby supporting quicker recovery and better overall maternal health.

Authors Contribution

Conceptualization: AA^1 Methodology: AA^1 , AA^2

Formal analysis: SA

Writing review and editing: SA, SN, AK, NA

All authors have read and agreed to the published version of the manuscript

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Koziol KJ, Smiley A, Latifi R, Castaldi MT. Predictive Risk Factors for Childbirth-Associated Breast Infections in the United States: A 10-Year Perspective. International Journal of Environmental Research and Public Health.2023Jul;20(14):6333.doi:10.3390/ijerph 20146333.
- [2] Lam E, Chan T, Wiseman SM. Breast Abscess: Evidence Based Management Recommendations. Expert Review of Anti-Infective Therapy.2014Jul;12 (7):753-62.doi:10.1586/14787210.2014.913982.

- [3] Kataria K, Srivastava A, Dhar A. Management of Lactational Mastitis and Breast Abscesses: Review of Current Knowledge and Practice.Indian Journal of Surgery.2013Dec;75:430-5.doi:10.1007/s12262-012-0776-1.
- [4] Algammal AM, Hetta HF, Elkelish A, Alkhalifah DH, Hozzein WN, Batiha GE et al. Methicillin-Resistant Staphylococcus Aureus (MRSA): One Health Perspective Approach to the Bacterium Epidemiology ,Virulence Factors, Antibiotic-Resistance, and Zoonotic Impact. Infection and Drug Resistance.2020 Sep: 3255-65. doi: 10.2147/IDR.S272733.
- [5] Sullivan T and de Barra E. Diagnosis and Management of Cellulitis.Clinical Medicine.2018Apr;18(2):160-3.doi: 10.7861/clinmedicine.18-2-160.
- [6] Chen C, Luo LB, Gao D, Qu R, Guo YM, Huo JL et al. Surgical Drainage of Lactational Breast Abscess with Ultrasound-Guided Encor Vacuum-Assisted Breast Biopsy System. The Breast Journal. 2019Sep;25(5):8 89-97.doi: 10.1111/tbj.13350.
- [7] Pal B, Shaikh O, Vijayakumar C, Prakash S, Balasubramanian G, Kumbhar U. A Randomized Controlled Trial Comparing Ultrasonography-Guided Needle Aspiration and Surgical Drainage for the Management of Breast Abscess.Cureus.2023Dec;15 (12).doi:10.7759/cureus.50956.
- [8] Irusen H, Rohwer AC, Steyn DW, Young T. Treatments for Breast Abscesses in Breastfeeding Women. Cochrane Database of Systematic Reviews.2015(8). doi:10.1002/14651858.CD010490.pub2.
- [9] Totadri VM, Vetri R, Sainath S. A Comparative Study of Drainage of Breast Abscesses by Conventional Incision and Drainage vs Ultrasound-Guided Needle Aspiration/Re-Aspiration in A Tertiary Health Care Centre.European Journal of Breast Health.2024Jul; 20(3):194. doi: 10.4274/ejbh.galenos.2024.2024-3-2.
- [10] Chandika AB, Gakwaya AM, Kiguli-Malwadde E, Chalya PL. Ultrasound Guided Needle Aspiration Versus Surgical Drainage in the Management of Breast Abscesses: A Ugandan Experience.BioMed Central Research Notes.2012 Dec;5:1-7.doi:10.1186/1756-050 0-5-12.
- [11] Afzal S, Bashir A, Shahzad H, Masroor I, Sattar AK. Ultrasound-Guided Percutaneous Aspiration for the Treatment of Breast Abscess at A Tertiary Care Center in the Developing World.Cureus.20220ct;14(10).doi: 10.7759/cureus.30865.
- [12] Boakes E, Woods A, Johnson N, Kadoglou N. Breast Infection: A Review of Diagnosis and Management Practices.European Journal of Breast Health.2018 Jul ;14(3): 136.
- [13] Singh SP, Kumar A, Chaudhary SK, Kumaran R. Comparison of Conventional Incision and Drainage Versus Ultrasound Guided Minimally Invasive Techniques in Management of the Breast Abscess: A Prospective Cohort Study.International Journal of Health Sciences.2022;6(S2):4349-57.doi:10.53730/ ijhs.v6nS2.6001.
- [14] Hussain N, Khan I, Ahmed T, Parveen S, Malik M, Khan MI. Comparison of the Restoration of Breast Feeding

DOI: https://doi.org/10.54393/pjhs.v6i5.1725

After Percutaneous Aspiration Vs Incision and Drainage for Management of Breast Abscess. Journal of Liaquat University of Medical and Health Sciences. 2018; 17(01): 47-51. doi: 10.22442/jlumhs.181710548.

- [15] Karim SA, Sami FA, Shah AH, Hanif MU, Zai AR. Compare the Outcome of Percutaneous Aspiration with Incisional Drainage for Management of Breast Abscess.Pakistan Journal of the Medical Health Science.2021;15(8):2157-9.doi:10.53350/pjmhs211582 157.
- [16] Manzoor A, Rashid I, Waqar SH, Shah SA. Comparison of Needle Aspiration and Incision and Drainage in the Management of Lactational Breast Abscess. Journal of The Society of Obstetricians and Gynaecologists of Pakistan.2022Mar;12(1):32-7.
- [17] Voruganti MR, Sistu MK, Gadipudi HC. A Comparative Study on Ultrasound Guided Percutaneous Aspiration Versus Incision and Drainage in the Management of Small Breast Abscesses.International Surgery Journal.2022Feb;9(3):620-6.doi:10.18203/2349-2902 .isj20220632.
- [18] Randhawa SR, Akram M, Akram H, Sajid M. Comparison of Needle Aspirations and Incision and Drainage of Breast Abscess. Journal of University Medical and Dental College.2019 Jun; 10(2): 40-4.
- [19] Zhou F, Li Z, Liu L, Wang F, Yu L, Xiang Y et al. The Effectiveness of Needle Aspiration Versus Traditional Incision and Drainage in the Treatment of Breast Abscess: A Meta-Analysis.Annals of Medicine.2023 Dec;55(1):2224045.doi:10.1080/07853890.2023.222 4045.
- [20] Bing F and Jie L. Ultrasound Guided Needle Aspiration and Cavity Washing Versus Incision and Drainage to Treat Breast Abscesses: A Meta-Analysis. International Journal of Clinical and Experimental Medicine. 2017 Jan; 10(6): 8656-65.