



Original Article



Association of Physical Activity With Perceived Stress and Well-Being in the Third Trimester of Pregnancy

Waqar Ahmed^{1*}, Parveen Akhter¹, Fatima Soomro¹, Husan Bano Channar¹, Waheed Ahmed², Saad Raza¹ and Abdul Qudoos³¹People's Nursing School, Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan²Dow University of Health Sciences, Karachi, Pakistan³Allama Iqbal Open University, Islamabad, Pakistan

ARTICLE INFO

Keywords:

Physical Activity, Perceived Stress, Prenatal Care, Third Trimester of Pregnancy

How to Cite:Ahmed, W., Akhter, P., Soomro, F., Channar, H. B., Ahmed, W., Raza, S., & Qudoos, A. (2025). Association of Physical Activity With Perceived Stress and Well-Being in the Third Trimester of Pregnancy: Physical Activity in the Third Trimester of Pregnancy. *Pakistan Journal of Health Sciences*, 6(1), 179-184. <https://doi.org/10.54393/pjhs.v6i1.2636>***Corresponding Author:**

Waqar Ahmed

People's Nursing School, Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan
waqar.ahmed@my.lumhs.edu.pkReceived date: 5th December, 2024Acceptance date: 23rd January, 2025Published date: 31st January, 2025

ABSTRACT

Physical inactivity is considered a worldwide pandemic that leads to numerous health problems. The World Health Organization advises pregnant women to participate in at least 150 minutes of moderate exercise each week to enhance health outcomes. As in the last trimester of pregnancy, considerable physical, hormonal, and psychological changes at this stage need further exploration. **Objectives:** To determine the association between physical activity, perceived stress, and well-being in the third trimester of pregnancy. **Methods:** A cross-sectional study of 245 third-trimester pregnant mothers was conducted at a local healthcare center using non-probability purposive sampling from June 2024 to November 2024. The Pregnancy Physical Activity Questionnaire (PPAQ) was used to quantify physical activity, stress levels were measured with the Perceived Stress Scale-04 (PSS-04), and well-being was assessed using the WHO-5 Well-Being Index. Spearman correlation analysis and descriptive statistics were used to investigate the connections among stress, physical activity, and well-being. **Results:** Increased physical activity is associated with decreased levels of stress, based on the data, It demonstrated a strong inverse relationship between physical activity and perceived stress ($\rho=-0.342$, $p<0.01$). Additionally, there was a positive correlation between well-being and physical exercise ($\rho=0.232$, $p<0.01$). **Conclusions:** It was concluded that physical activity decreased stress and improved well-being in the third trimester of pregnant women. This implies that medical professionals should recommend physical activities in prenatal care, especially in various cultural needs programs. More research is needed with larger, diverse groups.

INTRODUCTION

Physical inactivity has become a global public health issue, with over 80% of adolescents and 31% of adults not reaching the World Health Organization's (WHO) recommended levels of physical activity. Recently, physical inactivity has been recognized as the fourth leading contributor to premature deaths worldwide, escalating to pandemic proportions [1, 2]. This widespread inactivity has been linked to several chronic health problems, such as obesity, diabetes, and cardiovascular diseases, as well as a rise in psychological disorders, such as depression, anxiety, and stress [3]. One effective way to prevent these chronic diseases is by increasing physical activity. Encouraging active lifestyles addresses the root causes of

many emerging chronic conditions. This strategy supports "more active people for a healthier world" as outlined in the WHO's Global Action Plan on Physical Activity (2018–2030) [4]. In recent times, most developed nations have changed their policies toward maternal care and the prevention of ailments rather than treatment. In this regard, the approach is to catch the illness before it develops; likewise, in pregnancy, everything is related to the mom and the kid. Furthermore, early development of health behaviour will help reduce a specific chronic disease [7]. Pregnancy, especially during the third trimester, is a transformative period marked by significant physical, emotional, psychological, and hormonal changes as the body prepares



for childbirth. Hormonal fluctuations during this stage can affect mood, energy levels, and overall well-being. These changes, coupled with physical discomfort and concerns about childbirth, can lead to heightened stress and anxiety. These factors highlight the importance of understanding and addressing the unique needs of pregnant women during this critical phase [6]. Mental well-being during pregnancy is important because high levels of stress are known to affect not just the mother but also her unborn child. Pregnancy stress affects not only the emotional and physical health of the pregnant mother but also the growth and development of the unborn baby, contributing to low birth weight, preterm delivery, and developmental delays. It is, therefore, very important to ensure that mental health is considered during pregnancy for the well-being of both mother and child [7, 8]. Studies suggest that regular physical activity can play a significant role in preventing prenatal depression by improving mood and alleviating stress [9]. To improve health outcomes, the WHO advises pregnant mothers to engage in at least 150 minutes of moderate exercise each week [10]. However, approximately one in five pregnant women continue to experience suicidal thoughts, primarily due to untreated or severe depression, underscoring the need for effective interventions to address mental health challenges during pregnancy [11]. While global research supports the benefits of physical activity in stimulating the release of mood-enhancing hormones like serotonin [12], studies focusing on its impact during the third trimester when hormonal changes are most pronounced are limited, particularly in Pakistan. This gap in research highlights the importance of investigating the relationship of physical activity with perceived stress, also well-being during this critical period in Pakistan's unique cultural context. This research aims to create a healthier population by promoting increased physical activity, ultimately reducing the global burden of chronic diseases. Public health initiatives that encourage regular exercise strive to lower the rates of conditions like diabetes, heart disease, and obesity, thereby improving overall quality of life and reducing healthcare costs worldwide [13]. These efforts also support achieving Sustainable Development Goal 3 (SDG 3), which aims to promote well-being and healthy lives [14]. The findings from this study aim to provide valuable insights for healthcare organizations and policymakers in Pakistan, enabling them to develop strategies that integrate physical activity promotion into prenatal care programs. This research contributes to global efforts to address health disparities, improve maternal health outcomes, and foster healthier, more active populations, ultimately supporting sustainable development and a better quality of life. Physical inactivity is a global issue linked to poor health outcomes [15]. The WHO

recommends 150 minutes of moderate exercise weekly [10], but cultural norms in Pakistan often discourage activity during pregnancy [16]. Addressing this gap can improve health outcomes and align with global health goals like SDG 3.

This study aims to assess physical activity levels during the third trimester of pregnancy and investigate their relationship with perceived stress and well-being.

METHODS

An analytical cross-sectional study was conducted at the Gynecology and Obstetrics Outpatient Department (OPD) of Liaquat University Hospital, Hyderabad, to explore the association of physical activity with stress and well-being in the third trimester of pregnancy. The study was conducted over six months, from June 2024 to November 2024, after obtaining approval from the Ethical Review Committee (ERC) of Liaquat University of Medical and Health Sciences (LUMHS) (Approval number: LUMHS/REC/303). A sample size of 245 was calculated using OpenEpi (prevalence: 17.6%, confidence level: 95%, margin of error: 5%), with an additional 10% included to account for non-responses or dropouts. Pregnant women in their third trimester were selected using non-probability purposive sampling. Participants included women aged 18–35 years attending the Gynecology OPD who provided informed consent and were free from severe medical or obstetric complications affecting physical activity, stress, or well-being. Women were excluded if they did not consent, were advised against physical activity for medical reasons, had physical disabilities, or were on psychological medications. Data were collected using structured questionnaires, which were freely available, validated, and reliable. Minor modifications were made to adapt to the local context, and experts reviewed the validation of these modifications. The demographic details section collected participant information, including age, residence, education, and other pertinent variables. The Pregnancy Physical Activity Questionnaire (PPAQ) assessed daily physical activity, with higher scores indicating greater activity levels. This tool has shown strong reliability, with a Cronbach's alpha greater than 0.7. The Perceived Stress Scale (PSS-04) evaluated stress levels using four items rated on a 0–4 Likert scale, where higher scores correspond to increased stress. This scale also shows good reliability (Cronbach's alpha >0.7). The WHO-5 Well-Being Index also measured overall well-being through five items rated on a 0–5 scale, with higher scores reflecting better well-being. This index has a high reliability level, with a Cronbach's alpha of more than 0.8. To analyze the data, SPSS version 22.0 was used. Descriptive statistics, such as percentages and frequencies, were computed for categorical variables. Correlation analysis between physical activity, stress, and

well-being was conducted using the Pearson correlation test, with confidence intervals added for comprehensive reporting. At a 95% confidence level, a p-value of less than 0.05 was deemed statistically significant. Ethical considerations included obtaining written informed consent, ensuring participant confidentiality, and adhering to ERC guidelines.

RESULTS

A total of 245 women in their third trimester participated in the study. The sample's demographic characteristics offer a meaningful context for understanding the study findings. The average age of the participants was 25.05 ± 4.60 years, with an age range extending from 18 to 35 years. Most of the participants were between the ages of 18 and 30 years, which is reflective of the typical reproductive age range in this population. This data suggests that most participants were Urdu-speaking (76.3%), followed by Sindhi (20.0%) and others (3.7%). Additionally, 48.2% of participants had education above the secondary level, while 29.4% had no formal education. These demographics may have influenced the findings, which are interrelated with physical activity, well-being, and stress during pregnancy (Table 1).

Table 1: Sociodemographic Characteristics of Participants

Variable	Categories	Frequency (%)
Age group (Years)	18-25	146 (59.6)
	26-30	78 (31.8)
	31-35	21 (8.6)
Ethnicity	Urdu	187 (76.3)
	Sindhi	49 (20.0)
	Other (Saraiki, Punjabi)	9 (3.7)
Education Level	None	72 (29.4)
	Primary	55 (22.4)
	Secondary and Above	118 (48.2)

The analysis of physical activity levels among participants revealed that most engaged in light physical activity, contributing 69% (9,767 METs), followed by moderate activity at 20% (2,806 METs). Sedentary behaviour accounted for 11% (1,564 METs), with no participants reporting vigorous physical activity. These findings underscore a preference for low-intensity activities during pregnancy, highlighting the need for interventions to encourage moderate-intensity activities for better maternal health outcomes (Figure 1).




Figure 1: Cumulative Physical Activity Distribution Across Participants

Stress levels were assessed among participants, revealing that 63.7% experienced normal stress levels, while 36.3% exhibited elevated stress levels. This finding emphasizes the need for targeted interventions to alleviate stress, especially during the third trimester when pregnancy's physical and emotional demands peak (Figure 2).




Figure 2: Distribution of Stress Levels Among Participants

Mental well-being analysis indicated that 21.2% of pregnant women experienced depression, 31.8% reported poor well-being, and 46.9% had better well-being. These results underscore the mental health challenges prevalent during pregnancy, particularly in the third trimester, highlighting the need for focused interventions (Figure 3).




Figure 3: Distribution of mental well-being Levels Among Participants

Spearman's correlation analysis revealed a negative and moderate relation between perceived stress with physical activity engagement ($r = -0.342$, $p < 0.01$), which means high physical activity levels are associated with lower perceived stress (Table 2).

Table 2: Spearman's Correlation Analysis Between Physical Activity and Perceived Stress

Variable	Physical Activity	Perceived Stress
Physical Activity	1.000	-0.342 (p<0.01)
Perceived Stress	-0.342 (p<0.01)	1.000

Similarly, a weak but significant positive correlation was found between physical activity and well-being ($r=0.232$, $p<0.01$). This indicates that increased physical activity is mildly associated with improved well-being (Table 3).

Table 3: Spearman's Correlation Analysis Between Physical Activity and Well-Being

Variable	Physical Activity	Well-Being
Physical Activity	1.000	0.232 (p<0.01)
Well-Being	0.232 (p<0.01)	1.000

DISCUSSION

The sample's demographic characteristics offer a meaningful context for understanding the study findings. The average age of the mothers was 25.05 ± 4.60 years, with the age range extending from 18 to 35 years as shown in Table 1, reflecting the typical reproductive age range. Younger participants, generally more active than older women, may have influenced physical activity and psychological health findings. The relatively young age of the participants is important to note as it may influence both physical activity and psychological health, given that younger women generally engage in more physical activity than older women [17]. This demographic information helps contextualize the physical activity behaviours observed in the study. In this study, the majority of physical activity was light, accounting for 69% of the total activity. Moderate physical activity constituted 20%, while sedentary activity made up 11%. Notably, no participants reported engaging in vigorous physical activity. This distribution aligns with previous research, which shows that pregnant women mostly engaged in light to moderate-intensity physical activities [18]. This study found no evidence of vigorous physical activity during the third trimester. Similarly, a North Carolina study on pregnant women reported that vigorous activity was uncommon during pregnancy [19]. Results showed that 63.7% of participants had normal stress, while 36.3% experienced elevated levels. Elevated stress during pregnancy is linked to risks like preterm births and low birth weights. Elements like hormonal changes, social support, and discomfort can increase stress, consistent with findings from previous studies [20]. Well-being was measured using the WHO-5 index. A score above 50 indicated better well-being, 50 or below indicated poor well-being, and below 28 signified depression. The total score, scaled to 100, was calculated by multiplying the raw score by 4, as defined in previous studies [21]. 21.2% of pregnant women experienced

depression, while 31.8% reported poor well-being, indicating common mental health challenges during the third trimester. However, 46.9% maintained better well-being, highlighting resilience in many mothers. These findings align with WHO reports that nearly 1 in 5 women, or 20%, face mental health issues during pregnancy or postpartum, including suicidal thoughts or self-harm. Neglecting maternal mental health can negatively impact both the mother's and the infant's development [22]. The results underline the connection between perceived stress and physical activity, indicating that some lower stress levels while pregnant may be a result of increased physical activity. The Spearman correlation analysis supports this observation, showing a moderate negative relationship between perceived stress and physical activity ($r=-0.342$, $p<0.01$), showing that being more active is linked to lower levels of perceived stress, these results align with existing evidence emphasizing the stress-reducing benefits of physical activity. Being more active is known to stimulate the release of endorphins, which can improve mood and reduce stress hormones like cortisol [23]. Additionally, The Spearman correlation analysis reveals a significant positive relationship between physical activity and well-being ($r=0.232$, $p<0.01$). Although the correlation is weak, this could be because well-being is influenced by other factors such as social support, personal circumstances, and access to healthcare [24]. Despite the modest strength of the correlation, the results suggest that physical activity can play a role in enhancing well-being. These findings are consistent with research showing that physical activity helps reduce stress and contributes to better overall well-being. Encouraging pregnant women to engage in regular physical activity, even at light or moderate levels, can promote mental and physical health during pregnancy [25, 26]. The study emphasizes the importance of promoting safe, moderate physical activity during pregnancy to reduce stress and improve well-being. Culturally sensitive programs, community involvement, and stress-reduction strategies like counselling and mindfulness should be integrated into prenatal care. Enhanced mental health services and family support are also essential. Further research is needed to explore how ethnicity influences physical activity, stress, depression, and well-being during pregnancy.

CONCLUSIONS

It was concluded that a significant positive correlation was found between physical activity and well-being, and a significant negative correlation was found between physical activity and perceived stress, suggesting that increased activity is slightly associated with better well-being and reduced stress. It was also found that most mothers were performing light and fewer mothers were

performing moderate physical activity, which the World Health Organization recommends. These findings emphasize the importance of promoting physical activity during pregnancy for improved mental health and overall well-being.

Authors Contribution

Conceptualization: WA¹

Methodology: WA¹, PA, AQ

Formal analysis: WA¹, PA, FS, HBC, WA², SR, AQ

Writing review and editing: WA¹, PA, FS, HBC, WA², SR

All authors have read and agreed to the published version of the manuscript

Conflicts of Interest

The authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Guthold R, Stevens GA, Riley LM, Bull FC. Global Trends in Insufficient Physical Activity Among Adolescents: A Pooled Analysis of 298 Population-Based Surveys With 1.6 Million Participants. *The Lancet Child and Adolescent Health*. 2020 Jan; 4(1): 23-35. doi: 10.1016/S2352-4642(19)30323-2.
- [2] Pratt M, Varela AR, Salvo D, Kohl III HW, Ding D. Attacking the Pandemic of Physical Inactivity: What Is Holding Us Back? *British Journal of Sports Medicine*. 2020 Jul; 54(13): 760-2. doi: 10.1136/bjsports-2019-101392.
- [3] Laar RA, Shi S, Ashraf MA, Khan MN, Bibi J, Liu Y. Impact of Physical Activity On Challenging Obesity in Pakistan: A Knowledge, Attitude, and Practice (KAP) Study. *International Journal of Environmental Research and Public Health*. 2020 Nov; 17(21): 7802. doi: 10.3390/ijerph17217802.
- [4] World Health Organization. Global Action Plan On Physical Activity 2018-2030: More Active People for a Healthier World. World Health Organization. 2019 Jan.
- [5] Babatunde AO, Shobanke HA, Akinade AA, Michael AJ, Osadare M, Akanbi OK et al. Enhancing Preventive Medicine Over Curative Medicine: Role of Telemedicine. *Public Health in Practice*. 2021 Nov; 2: 100130. doi: 10.1016/j.puhip.2021.100130.
- [6] Mah BL, Pringle KG, Weatherall L, Keogh L, Schumacher T, Eades S et al. Pregnancy Stress, Healthy Pregnancy and Birth Outcomes—The Need for Early Preventative Approaches in Pregnant Australian Indigenous Women: A Prospective Longitudinal Cohort Study. *Journal of Developmental Origins of Health and Disease*. 2019 Feb; 10(1): 31-8. doi: 10.1017/S204017441800079X.
- [7] Kołomańska-Bogucka D, Micek A, Mazur-Bialy AI. The COVID-19 Pandemic and Levels of Physical Activity in the Last Trimester, Life Satisfaction and Perceived Stress in Late Pregnancy and in the Early Puerperium. *International Journal of Environmental Research and Public Health*. 2022 Mar; 19(5): 3066. doi: 10.3390/ijerph19053066.
- [8] Wang W, Wen L, Zhang Y, Wang L, Wang L, Chen Z et al. Maternal Prenatal Stress and Its Effects On Primary Pregnancy Outcomes in Twin Pregnancies. *Journal of Psychosomatic Obstetrics and Gynecology*. 2020 Jul; 41(3): 198-204. doi: 10.1080/0167482X.2019.1611776.
- [9] Kołomańska-Bogucka D and Mazur-Bialy AI. Physical Activity and the Occurrence of Postnatal Depression—A Systematic Review. *Medicina*. 2019 Sep; 55(9): 560. doi: 10.3390/medicina55090560.
- [10] World Health Organization. Global Status Report On Physical Activity 2022: Country Profiles. World Health Organization. 2022 Dec.
- [11] Bhamani SS, Van Parys AS, Arthur D, Letourneau N, Wagnild G, Degomme O. Promoting Mental Wellbeing in Pregnant Women Living in Pakistan with the Safe Motherhood—Accessible Resilience Training (SM-ART) Intervention: A Randomized Controlled Trial. *BioMed Central Pregnancy and Childbirth*. 2024 Jun; 24(1): 452. doi: 10.1186/s12884-024-06629-2.
- [12] Dinas PC, Koutedakis Y, Flouris AD. Effects of Exercise and Physical Activity On Depression. *Irish Journal Of Medical Science*. 2011 Jun; 180: 319-25. doi: 10.1007/s11845-010-0633-9.
- [13] Chen S, Ma J, Hong J, Chen C, Yang Y, Yang Z et al. A Public Health Milestone: China Publishes New Physical Activity and Sedentary Behaviour Guidelines. *Journal of Activity, Sedentary and Sleep Behaviors*. 2022 Dec; 1(1): 9. doi: 10.1186/s44167-022-00009-x.
- [14] Das T, Holland P, Ahmed M, Husain L, Ahmed M, Husain L. Sustainable Development Goal 3: Good Health and Well-Being. In *South-East Asia Eye Health: Systems, Practices, and Challenges*. Singapore: Springer Singapore. 2021 Aug; 61-78. doi: 10.1007/978-981-16-3787-2_4.
- [15] Musa S, Dergaa I, Bachiller V, Saad HB. Global Implications Of COVID-19 Pandemic On Adults' Lifestyle Behavior: The Invisible Pandemic of Non-Communicable Disease. *International Journal of Preventive Medicine*. 2023 Feb; 14(1): 15. doi: 10.4103/ijpvm.ijpvm_157_21.
- [16] Yaseen M. The Knowledge, Attitude and Practices of Pregnant Women Regarding Physical Activity in

- Turbat, Kech. *Annals of Social Sciences and Perspective*. 2022 Dec; 3(2): 297-309. doi: 10.52700/assap.v3i2.176.
- [17] Nemoto Y, Brown WJ, Mielke GI. Trajectories of Physical Activity from Mid to Older Age in Women: 21 Years of Data from the Australian Longitudinal Study on Women's Health. *International Journal of Behavioral Nutrition and Physical Activity*. 2024 Jan; 21(1): 4. doi: 10.1186/s12966-023-01540-z.
- [18] Nadeem S, Khatoon A, Rasheed S, Munim TF. The Physical Activity Patterns Among Pregnant Women at A Tertiary Care Hospital in, Pakistan. *Pakistan Journal of Medical Sciences*. 2022 Mar; 38(4Part-II): 904. doi: 10.12669/pjms.38.4.4809.
- [19] Gascoigne EL, Webster CM, Honart AW, Wang P, Smith-Ryan A, Manuck TA. Physical Activity and Pregnancy Outcomes: An Expert Review. *American Journal of Obstetrics and Gynecology Maternal Fetal Medicine*. 2023 Jan; 5(1): 100758. doi: 10.1016/j.ajogmf.2022.100758.
- [20] Răchită A, Strete GE, Suciuc LM, Ghiga DV, Sălcudean A, Mărginean C. Psychological Stress Perceived By Pregnant Women in the Last Trimester of Pregnancy. *International Journal of Environmental Research and Public Health*. 2022 Jul; 19(14): 8315. doi: 10.3390/ijerph19148315.
- [21] Omani-Samani R, Maroufizadeh S, Almasi-Hashiani A, Sepidarkish M, Amini P. The WHO-5 Well-Being Index: A Validation Study in People With Infertility. *Iranian Journal of Public Health*. 2019 Nov; 48(11): 2058. doi: 10.18502/ijph.v48i11.3525.
- [22] Yu H, Shen Q, Bränn E, Yang Y, Oberg AS, Valdimarsdóttir UA et al. Perinatal Depression and Risk of Suicidal Behavior. *Journal of American and Medical Association Network Open*. 2024 Jan; 7(1): e2350897-. doi: 10.1001/jamanetworkopen.2023.50897.
- [23] Cao B, Zhao Y, Ren Z, McIntyre RS, Teopiz KM, Gao X et al. Are Physical Activities Associated with Perceived Stress? The Evidence from the China Health and Nutrition Survey. *Frontiers in Public Health*. 2021 Aug; 9: 697484. doi: 10.3389/fpubh.2021.697484.
- [24] Evanoff BA, Strickland JR, Dale AM, Hayibor L, Page E, Duncan JG et al. Work-Related and Personal Factors Associated with Mental Well-Being During the COVID-19 Response: Survey of Health Care and Other Workers. *Journal Of Medical Internet Research*. 2020 Aug; 22(8): e21366. doi: 10.2196/21366.
- [25] Sánchez-Polán M, Silva-Jose C, Franco E, Nagpal TS, Gil-Ares J, Lili Q et al. Prenatal Anxiety and Exercise. Systematic Review and Meta-Analysis. *Journal of Clinical Medicine*. 2021 Nov; 10(23): 5501. doi: 10.3390/jcm10235501.
- [26] Guinhouya BC, Duclos M, Enea C, Storme L. Beneficial Effects of Maternal Physical Activity During Pregnancy On Fetal, Newborn, and Child Health: Guidelines for Interventions During the Perinatal Period from the French National College of Midwives. *Journal of Midwifery and Women's Health*. 2022 Nov; 67: S149-57. doi: 10.1111/jmwh.13424.