



Original Article



Evaluating the Incidence of Co-Existing Injuries in Anterior Talofibular Ligament Injuries a Magnetic Resonance Imaging Study

Farhan Qazi¹, Ojala Shaheen², Wajeeha Imran Andrabi³, Muhammad Arif⁴, Farzana Begum⁵ and Mahwash Mansoor⁶

¹Department of Orthopedics, Pak International Medical College, Peshawar, Pakistan

²Department of Radiology, Islamabad Diagnostic Center, Islamabad, Pakistan

³Department of Radiology, Lahore Medical and Dental College, Lahore, Pakistan

⁴Department of Orthopedics, Akhtar Saeed Medical College, Rawalpindi, Pakistan

⁵Department of Radiology, Pak International Medical College, Peshawar, Pakistan

⁶Department of Radiology, Bolan Medical College, Quetta, Pakistan

ARTICLE INFO

Keywords:

Anterior Talofibular Ligament, Calcaneofibular Ligament, Ankle Injuries, Osteochondral Defect

How to Cite:

Qazi, F., Shaheen, O., Andrabi, W. I., Arif, M., Begum, F., & Mansoor, M. (2025). Evaluating the Incidence of Co-Existing Injuries in Anterior Talofibular Ligament Injuries a Magnetic Resonance Imaging Study: Co-Existing Injuries in Anterior Talofibular Ligament . Pakistan Journal of Health Sciences, 6(3), 115-120. <https://doi.org/10.54393/pjhs.v6i3.2667>

***Corresponding Author:**

Farhan Qazi
 Department of Orthopedics, Pak International Medical College, Peshawar, Pakistan
dr_farhanqazi@yahoo.com

Received date: 29th November, 2024

Acceptance date: 14th March, 2025

Published date: 31st March, 2025

ABSTRACT

Ankle lateral ligament injuries are common in everyday life as well as in athletic activities. Ankle injuries most commonly affect the anterior talofibular ligament (ATFL). Due to ATFL's susceptibility, achieving intelligent localization and injury evaluation is extremely important from a clinical standpoint. **Objectives:** To evaluate the incidence and patterns of co-existing injuries in ATFL damage using Magnetic Resonance Imaging (MRI). **Methods:** In this cross-sectional study, fifty-five patients undergoing surgical management for ATFL injuries were included. Preoperative MRIs of affected ankles were analyzed to document ATFL integrity and associated injuries. Descriptive statistics were used to summarize findings, with categorical variables reported as frequencies and percentages and continuous variables as means \pm SD. **Results:** The mean age of the patients was 36.8 ± 12.4 years, with 61.8% male and 38.2% female. Associated injuries were observed in 72.7% of patients. ATFL injuries included 30.9% low-grade incomplete, 21.8% high-grade incomplete, and 14.5% complete tears. calcaneofibular ligament (CFL) injuries were common, with 36.4% showing partial tears, and 3.6% complete tears. The deep deltoid ligament was intact in 56.4%, while 43.6% had injuries. The superficial deltoid ligament was intact in 70.9%. The peroneal tendon was intact in 83.6%, with 5.5% having injuries and 10.9% showing tendinosis. Osteochondral Defects (OCDs) were present in 21.8% of cases. **Conclusions:** It was concluded that ATFL injuries frequently occur with damage to other structures, particularly the CFL and deltoid ligaments. MRI facilitates accurate diagnosis, emphasizing the need for comprehensive assessment and concurrent management of associated injuries during ATFL repair.

INTRODUCTION

In sports, Ankle sprains are among the most common injuries, comprising up to 40% of all athletic injuries [1]. Despite their frequency, the patterns of associated structural damage, optimal diagnostic approaches, and their clinical significance continue to be areas of clinical interest. The ankle joint, which is formed by the tibia, fibula, and talus bones, is a hinged synovial joint. This joint is supported by the medial and lateral ligament complexes as well as the syndesmotic ligaments. Among these, the lateral ligament complex is most vulnerable to injury,

particularly during inversion movements. These movements account for the majority of ankle sprains. This ligament complex includes the ATFL, the calcaneofibular ligament (CFL), and the posterior talofibular ligament (PTFL). The ATFL is the most frequently injured structure due to its primary role in stabilizing the ankle during plantarflexion and resisting anterolateral talar translation. Studies suggest that the ATFL is damaged in as many as 90% of significant ankle injuries, compared to 50-75% for the CFL and only 10% for the PTFL [2, 3]. Lateral ankle



sprains typically occur when excessive abduction and internal rotation of the rear foot are combined with external rotation of the lower leg. Such movements place undue tension on the lateral ligaments, particularly during plantarflexion, increasing the risk of ligamentous injury. Damage occurs when the tensile forces acting on these ligaments exceed their physiological strength [4, 5]. While the primary injury involves the ATFL, co-existing injuries to adjacent structures such as the CFL, deltoid ligaments, peroneal tendons, and osteochondral surfaces are frequently observed. These injuries can influence treatment decisions and long-term outcomes, making their accurate detection clinically significant. The initial clinical evaluation of an ankle sprain focuses on identifying dislocations or asymmetries. It also focuses on palpating for tenderness over the medial and lateral ankle ligaments, and assessing the fibula. To rule out more severe injuries, observations for edema, ecchymosis, muscle strength deficits, and range of motion restrictions are crucial, along with neurovascular examinations [6, 7]. Advanced imaging modalities, such as ultrasonography (US), MRI, arthrography, and stress radiography, play a key role in diagnosing ligament injuries, especially in chronic cases [8]. Although stress radiography is particularly useful in ruling out fractures, which occur in less than 15% of cases, US and MRI are more sensitive in detecting ligamentous injuries [9]. US is a valuable bedside tool, with a reported 91% accuracy in identifying ATFL injuries, while MRI demonstrates 97% accuracy, particularly for detecting associated injuries such as bone marrow edema, tendon abnormalities, and osteochondral defects [10]. However, the diagnostic sensitivity of MRI for acute ATFL injuries varies widely, ranging from 40% to 95%, with specificity reported between 70% and 97% [11, 12]. MRI findings for ATFL injuries may include ligament discontinuity, irregular contours, thickening or thinning, the bright rim sign [13], or bone avulsions [14, 15]. Despite these advantages, MRI is often underutilized in the acute setting due to its variable sensitivity, cost, and limited availability. Additionally, most existing literature focuses on chronic instability rather than the early detection of concurrent injuries in acute ATFL tears [16]. Persistent symptoms such as instability, stiffness, edema, pain, and muscle weakness are reported in 10%-30% of patients following initial treatment, underscoring the need for careful long-term management [12, 15]. While physical therapy is effective for most cases, surgical intervention may be required in patients with unresolved symptoms or chronic instability. A key limitation in the current understanding of ATFL injuries is the lack of studies assessing the prevalence and patterns of associated injuries using MRI in the acute phase. Identifying these injuries early can help tailor rehabilitation strategies and prevent long-term complications.

This study aims to systematically evaluate the prevalence and patterns of associated injuries in patients with ATFL tears using MRI. By identifying the structures most commonly affected alongside the ATFL, this research seeks to bridge a critical knowledge gap and improve both diagnostic accuracy and patient management strategies.

METHODS

A cross-sectional study was conducted over six months commencing from March, 2024, up till August, 2024. This study was conducted at Pak International Medical College, Hayatabad, Peshawar, Pakistan. A total of 55 patients were selected for the study, who experienced surgical procedures for ATFL injuries. Written informed consent was obtained from all participants. The Institutional Review Board (IRB) of Pak International Medical College has been permitted to carry out this study with reference number PIMC/DMR/2. The sample size of 55 was calculated using G*Power software, based on an estimated effect size of 0.5, a power of 80%, and a significance level (α) of 0.05. The effect size was determined from prior studies assessing associated injuries in ATFL tears. To minimize potential biases due to loss of follow-up, all participants were assessed at a single time point using MRI before surgery, ensuring a standardized evaluation of associated injuries. The sample was selected through purposive sampling. The sample focused on patients who fulfilled the defined inclusion and exclusion criteria. The participants undergoing a preoperative MRI scan of the affected ankle, with reports specifically addressing the integrity of the ATFL, deltoid ligaments, peroneal tendons, CFL, and the presence of an OCD were included. Patients with systemic diseases or fracturing the same ankle were excluded. Additionally, individuals who did not undergo a preoperative MRI scan with the surgery for ATFL injuries were also excluded from the study. To standardize the classification of structural integrity all MRI scans were analyzed by three investigators using predefined terminology. These investigators included two musculoskeletal radiologists with over five years of experience in MRI interpretation and one orthopaedic surgeon with expertise in foot and ankle injuries. To minimize variability in reporting, terms such as "scarred" or "scarring" were interpreted as "sprain" for the determinations of the study. In cases of disagreement regarding the interpretation of MRI findings, the investigators convened to reach a consensus. Data were examined by incorporating SPSS version 26.0 (IBM Corp., Armonk, NY, USA). Descriptive statistics were used to summarize the findings. Continuous variables are reported as means and standard deviations (SD), which include such as age and body mass index (BMI). On the other hand, categorical variables are presented as frequencies and

percentages, comprising gender distribution, injury types, and ligament integrity.

RESULTS

The mean age of the patients was 36.8 ± 12.4 years, with 61.8% male and 38.2% female. The mean BMI was 28.43 ± 6.7 kg/m². The right ankle was affected in 54.5% of cases, while the left was affected in 45.5% (Table 1).

Table 1: Patient Demographics (n=55)

Measurements		n (%)
Age in Years	19-64 Years	36.8 ± 12.4
Gender	Male	34 (61.8%)
	Female	21 (38.2%)
Body Mass Index (kg/m ²)	19.2-38.9 Kg/M ²	28.43 ± 6.7
Operated Side	Right	30 (54.5%)
	Left	25 (45.5%)

The majority of cases (29.1%) exhibited low-grade incomplete injuries, followed by high-grade incomplete injuries (23.6%) and complete injuries (14.5%). A smaller proportion of participants had an undamaged ATFL (7.3%), while 25.5% had some form of injury without further classification (Table 2).

Table 2: Extent of Anterior Talofibular Ligament (ATFL) Injury

ATFL Injury Type	n (%)
Undamaged	4 (7.3%)
Injured	14 (25.5%)
Low-Grade Incomplete Injury	16 (29.1%)
High-Grade Incomplete Injury	13 (23.6%)
Complete Injury	8 (14.5%)

A total of 72.7% of patients had associated injuries, while 27.3% presented isolated ATFL injuries. 32.7% of patients had an undamaged ATFL, while 52.7% had some degree of incomplete injury (30.9% low-grade, 21.8% high-grade). Complete ATFL tears were observed in 14.5% of cases. Associated injuries predominantly involved the CFL and deltoid ligaments, highlighting their frequent co-involvement in ATFL damage (Table 3).

Table 3: Extent and Nature of Injuries to the ATFL

Measurement	n (%)
Degree of Injury	
Single Injury	15 (27.3%)
Related Injuries	40 (72.7%)
ATFL Integrity	
Undamaged	18 (32.7%)
Injured	29 (52.7%)
Low-Grade Incomplete Injury	17 (30.9%)
High-Grade Incomplete Injury	12 (21.8%)
Whole Tear	8 (14.5%)

The CFL was intact in 60% of patients. Partial tears (low- and high-grade combined) accounted for 36.4% (20% low-grade, 16.4% high-grade), while complete tears were uncommon, seen in 3.6% of patients. The deep deltoid

ligament was intact in 56.4% of patients, while 43.6% had some degree of injury (29.1% low-grade, 14.5% high-grade). The superficial deltoid ligament was intact in 70.9% of cases, with 29.1% showing some level of injury (14.5% low-grade, 14.5% high-grade). The peroneal tendon was intact in 83.6% of patients, while 5.5% had injuries, and 10.9% had tendinosis. OCD was associated with 21.8% of cases (Table 4).

Table 4: Associated Injuries

Measurement	n (%)
Calcaneofibular ligament (CFL)	
Undamaged	33 (60%)
Injured	22 (40%)
Low-Grade Incomplete Injury	11 (20%)
High-Grade Incomplete Injury	9 (16.4%)
Complete Injury	2 (3.6%)
Deep Deltoid Ligament	
Undamaged	31 (56.4%)
Injured	24 (43.6%)
Low-Grade Incomplete Injury	16 (29.1%)
High-Grade Incomplete Injury	8 (14.5%)
Superficial Deltoid Ligament	
Undamaged	39 (70.9%)
Injured	16 (29.1%)
Low-Grade Incomplete Injury	8 (14.5%)
High-Grade Incomplete Injury	8 (14.5%)
Peroneal tendon	
Undamaged	46 (83.6%)
Injured	3 (5.5%)
Tendinosis	6 (10.9%)
Osteochondral Defect (OCD)	
Associated OCD	12 (21.8%)
No Associated OCD	43 (78.2%)

DISCUSSION

Ankle sprains, particularly those involving the lateral ligament complex, remain one of the most common injuries in both athletic and general populations. Among these, the most frequently injured is the ATFL often in combination with other anatomical structures such as the deltoid ligament complex, CFL, osteochondral surfaces and peroneal tendons. Accurate diagnosis and understanding of co-existing injuries are crucial for effective management. It is also important to prevent long-term complications like instability or osteoarthritis. MRI plays a vital role in evaluating such injuries by offering detailed insights into both acute and chronic damage. This study investigates the spectrum and prevalence of co-existing injuries associated with ATFL damage. Hence, providing valuable data to inform clinical decision-making [17]. The most frequently injured structure associated with ATFL injuries was the CFL, with 41.8% involvement. This high prevalence underscores the importance of understanding

the biomechanical relationship between the ATFL and CFL. The CFL plays a pivotal part in maintaining lateral stabilization and resisting inversion forces [18]. Given the shared mechanism of injury in lateral ankle sprains, concurrent CFL damage is common. Despite this, MRI sensitivity for detecting CFL injuries varies significantly, with accuracies reported between 66% for partial tears and 88% for complete tears [19]. This diagnostic challenge necessitates a combination of imaging and clinical tests, such as the medial talar tilt stress test, to enhance diagnostic accuracy when MRI findings are inconclusive [20, 21]. Similarly, the deltoid ligament complex was frequently involved despite its primary role in stabilizing the medial ankle. This is likely due to compensatory stresses placed on the medial structures when the lateral stabilizers fail. Up to 15% of inversion injuries have been reported to involve the deltoid ligament, a finding consistent with our study [22]. Interestingly, while deltoid ligament injuries are classically linked to eversion mechanisms, up to 15% of inversion injuries may also involve this complex [23]. In this study, the deep deltoid ligament was intact in 56.4% of patients, with partial tears (low- and high-grade) and sprains observed in the remaining cases. Similarly, the superficial deltoid ligament was intact in 70.9% of cases, with a smaller proportion showing sprains or partial tears. Failure to recognize deltoid ligament injuries may contribute to residual instability despite surgical correction of ATFL tears, highlighting the need for a comprehensive medial and lateral ligament evaluation. The peroneal tendons are key stabilizers in lateral ankle movements. These were intact in the majority of patients. However, tendinosis was noted in 10.9% of cases, while tears were observed in 5.5%. These conclusions are consistent with earlier studies that emphasize the role of the peroneus longus tendon in stabilizing the lateral ankle during inversion injuries [24]. Despite their critical role, peroneal tendon injuries are frequently overlooked during routine evaluations [25]. Given their contribution to ankle stability, clinicians should assess peroneal tendon integrity in all cases of ATFL injury. Moreover, surgical management of ATFL injuries should address any co-existing peroneal tendon damage to optimize outcomes and prevent recurrent instability [26]. Osteochondral defects (OCDs), often resulting from direct trauma or repetitive micro-trauma, were observed in 21.8% of cases. These defects commonly occur during ankle sprains when forced talar rotation generates compression and shear forces on the talar cartilage [27]. Cadaveric studies have corroborated this mechanism, demonstrating similar cartilage injuries under experimental inversion stress [28]. Additionally, chronic lateral instability due to ATFL injury has been linked to the development of OCDs, where repetitive micro-trauma and altered joint

biomechanics exacerbate cartilage damage [29, 30]. The prevalence of OCDs in this study underscores the importance of evaluating talar cartilage integrity during the assessment of ATFL injuries, as timely intervention can mitigate long-term consequences like post-traumatic arthritis.

CONCLUSIONS

It was concluded that this study highlights a high prevalence of associated injuries in ATFL tears, particularly in the CFL (41.8%) and deltoid ligament complexes, reinforcing the interconnected nature of lateral and medial ankle stability. Additionally, osteochondral defects were present in 21.8% of cases, underscoring the long-term impact of ATFL injuries on joint health. Despite this, surgical treatment often focuses on ATFL repair alone, potentially overlooking concurrent injuries that contribute to persistent instability. The findings emphasize the need for a comprehensive assessment of ligamentous, tendinous, and osteochondral structures in ATFL injuries.

Authors Contribution

Conceptualization: FQ

Methodology: FQ, MA, FB

Formal analysis: OS, WIA

Writing review and editing: MM

All authors have read and agreed to the published version of the manuscript

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Park YG, Ha CW, Park YB, Na SE, Kim M, Kim TS *et al.* Is It Worth to Perform Initial Non-Operative Treatment for Patients with Acute ACL Injury? A Prospective Cohort Prognostic Study. *Knee Surgery and Related Research*. 2021 Dec; 33:1-9. doi:10.1186/s43019-021-00094-3.
- [2] Delahunt E, Bleakley CM, Bossard DS, Caulfield BM, Docherty CL, Doherty C *et al.* Clinical Assessment of Acute Lateral Ankle Sprain Injuries (ROAST): 2019 Consensus Statement and Recommendations of the International Ankle Consortium. *British Journal of Sports Medicine*. 2018 Oct; 52(20):1304-10. doi:10.1136/bjsports-2017-098885.
- [3] Pacheco J, Guerra-Pinto F, Araújo L, Flora M, Alçada R, Rocha T *et al.* Chronic Ankle Instability Has No Correlation with the Number of Ruptured Ligaments in Severe Anterolateral Sprain: A Systematic Review and Meta-Analysis. *Knee Surgery, Sports Traumatology,*

- Arthroscopy.2021Nov;29(11):3512-24.doi:10.1007/s00167-021-06610-y.
- [4] Mugno AT and Constant D. Recurrent Ankle Sprain. Treasure Island: Stat-Pearls Publishing. 2020 Aug.
- [5] Chen ET, McInnis KC, Borg-Stein J. Ankle Sprains: Evaluation, Rehabilitation, and Prevention. *Current Sports Medicine Reports*.2019Jun;18(6):217-23.doi:10.1249/JSR.0000000000000603.
- [6] Barini M, Zagaria D, Licandro D, Pansini S, Airoidi C, Leigh M et al. Magnetic Resonance Accuracy in the Diagnosis of Anterior Talo-Fibular Ligament Acute Injury:A Systematic Review and Meta-Analysis. *Diagnostics*.2021Sep;11(10):1782.doi:10.3390/diagnostics11101782.
- [7] Alves T, Dong Q, Jacobson J, Yablon C, Gandikota G. Normal and Injured Ankle Ligaments On Ultrasonography with Magnetic Resonance Imaging Correlation. *Journal of Ultrasound in Medicine*.2019 Feb;38(2):513-28.doi:10.1002/jum.14716.
- [8] Vuurberg G, Hoorntje A, Wink LM, Van Der Doelen BF, Van Den Bekerom MP, Dekker R et al. Diagnosis, Treatment and Prevention of Ankle Sprains: Update of an Evidence-Based Clinical Guideline. *British Journal of Sports Medicine*.2018Aug;52(15):956-.doi:10.1136/bjsports-2017-098106.
- [9] Noda Y, Horibe S, Hiramatsu K, Takao R, Fujita K. Quick and Simple Test to Evaluate Severity of Acute Lateral Ankle Sprain. *Asia-Pacific Journal of Sports Medicine, Arthroscopy, Rehabilitation and Technology*.2021Jul;25:30-4.doi:10.1016/j.asmart.2021.05.003.
- [10] Kellett JJ, Lovell GA, Eriksen DA, Sampson MJ. Diagnostic Imaging of Ankle Syndesmosis Injuries: A General Review. *Journal of Medical Imaging and Radiation Oncology*.2018Apr;62(2):159-68.doi:10.1111/1754-9485.12708.
- [11] Salat P, Le V, Veljkovic A, Cresswell ME. Imaging in Foot and Ankle Instability. *Foot and Ankle Clinics*. 2018Dec;23(4):499-522.doi:10.1016/j.fcl.2018.07.011.
- [12] Li HY, Li WL, Chen SY, Hua YH. Increased ATFL-PTFL Angle Could Be an Indirect MRI Sign in Diagnosis of Chronic ATFL Injury. *Knee Surgery, Sports Traumatology, Arthroscopy*.2020Jan;28:208-12.doi:10.1007/s00167-018-5252-4.
- [13] Kim M, Choi YS, Jeong MS, Park M, Chun TJ, Kim JS et al. Comprehensive Assessment of Ankle Syndesmosis Injury Using 3D Isotropic Turbo Spin-Echo Sequences: Diagnostic Performance Compared with That of Conventional and Oblique 3-T MRI. *American Journal of Roentgenology*.2017Apr;208(4):827-33.doi:10.214/AJR.16.16985.
- [14] Tao H, Hu Y, Qiao Y, Ma K, Yan X, Hua Y et al. T2-Mapping Evaluation of Early Cartilage Alteration of Talus for Chronic Lateral Ankle Instability with Isolated Anterior Talofibular Ligament Tear or Combined with Calcaneofibular Ligament Tear. *Journal of Magnetic Resonance Imaging*.2018Jan;47(1):69-77.doi:10.1002/jmri.25745.
- [15] Elkaim M, Thès A, Lopes R, Andrieu M, Cordier G, Molinier F et al. Agreement Between Arthroscopic and Imaging Study Findings in Chronic Anterior Talo-Fibular Ligament Injuries. *Orthopaedics and Traumatology: Surgery and Research*.2018Dec;104(8):S213-8.doi:10.1016/j.otsr.2018.09.008.
- [16] Glazebrook M. Percutaneous Ankle Reconstruction of Lateral. *Managing Instabilities of the Foot and Ankle, An issue of Foot and Ankle Clinics of North America*. 2018 Nov;23(4):581.doi:10.1016/j.fcl.2018.07.013.
- [17] Li L, Gollhofer A, Lohrer H, Dorn-Lange N, Bonsignore G, Gehring D. Function of Ankle Ligaments for Subtalar and Talocrural Joint Stability During an Inversion Movement-An in Vitro Study. *Journal of Foot and Ankle Research*.2019Dec;12:1-8.doi:10.1186/s13047-019-0330-5.
- [18] Vega J, Malagelada F, Dalmau-Pastor M. Arthroscopic All-Inside ATFL and CFL Repair Is Feasible and Provides Excellent Results in Patients with Chronic Ankle Instability. *Knee Surgery, Sports Traumatology, Arthroscopy*.2020Jan;28:116-23.doi:10.1007/s00167-019-05676-z.
- [19] Koh D, Chandrakumara D, King CK. Incidence of Injuries Associated with Anterior Talofibular Ligament Injury Based On the Reporting of Magnetic Resonance Imaging. *Cureus*.2023Jul;15(7).doi:10.7759/cureus.41738.
- [20] Cao S, Wang C, Ma X, Wang X, Huang J, Zhang C. Imaging Diagnosis for Chronic Lateral Ankle Ligament Injury: A Systemic Review with Meta-Analysis. *Journal of Orthopaedic Surgery and Research*.2018Dec;13:1-4.doi:10.1186/s13018-018-0811-4.
- [21] Larkins CG, Brady AW, Aman ZS, Dornan GJ, Haytmanek CT, Clanton TO. Evaluation of the Intact Anterior Talofibular and Calcaneofibular Ligaments, Injuries, and Repairs with and Without Augmentation: A Biomechanical Robotic Study. *The American Journal of Sports Medicine*.2021Jul;49(9):2432-8.doi:10.1177/03635465211018645.
- [22] Koris J, Calder JD, Dalmau-Pastor M, Fernandez MA, Ramasamy A. Deltoid Ligament Injuries: A Review of the Anatomy, Diagnosis and Treatments. *Knee Surgery, Sports Traumatology, Arthroscopy*.2024 Dec;32(12):3052-64.doi:10.1002/ksa.12274.
- [23] Gentile MA, Rigby RB, Fleming J, Taylor S, Lamba C. Chronic Medial Ankle Instability Medial Drive Through Sign and Surgical Repair Technique. *Foot & Ankle Surgery: Techniques, Reports and Cases*.2022Sep;2(3):100225.doi:10.1016/j.fastrc.2022.100225.
- [24] Mendez-Rebolledo G, Guzmán-Venegas R, Cruz-Montecinos C, Watanabe K, Calatayud J,

- Martinez-Valdes E. Individuals with Chronic Ankle Instability Show Altered Regional Activation of the Peroneus Longus Muscle During Ankle Eversion. *Scandinavian Journal of Medicine and Science in Sports*.2024Jan;34(1): e14535. doi: 10.1111/sms.14535.
- [25] Flor-de-Lima B, Castro M, Costa AS, Brito N, Madureira AJ. Magnetic Resonance Imaging of Common and Uncommon Peroneal Tendons Disorders. *Acta radiology*.2022 Apr; 34(1): 9-16.
- [26] Roch FE, Hahn FM, Jäckle K, Meier MP, Stinus H, Lehmann W et al. Diagnosis, Treatment, and Prevention of Ankle Sprains: Comparing Free Chatbot Recommendations with Clinical Guidelines. *Foot and Ankle Surgery*.2024Dec.doi:10.1016/j.fas.2024.12.003.
- [27] 27. Shimozone Y, Rammelt S, Takao M. Current Lack of Evidence On Treatment Strategies and Clinical Outcomes for Osteochondral Lesions of the Subtalar, Talonavicular, and Calcaneocuboid Joints:A Systematic Review. *Cartilage*.2024Mar;15(1):7-15.doi: 10.1177/19476035231216182.
- [28] Casado-Hernández I, Becerro-de-Bengoa-Vallejo R, Losa-Iglesias ME, Santiago-Nuño F, Mazo-terras-Pardo V, López-López D et al. Association between Anterior Talofibular Ligament Injury and Ankle Tendon, Ligament, and Joint Conditions Revealed by Magnetic Resonance Imaging. *Quantitative Imaging in Medicine and Surgery*.2021Jan;11(1):84.doi:10.21037/qims-20-5
- [29] Barbier O, Amouyel T, de l'Escalopier N, Cordier G, Baudrier N, Benoist J et al. Osteochondral Lesion of the Talus: What Are We Talking About? *Orthopaedics and Traumatology: Surgery and Research*.2021Dec; 107(8): 103068. doi: 10.1016/j.otsr.2021.103068.
- [30] Kim YS, Kim TY, Koh YG. Demographic Predictors of Concomitant Osteochondral Lesion of the Talus in Patients with Chronic Lateral Ankle Instability. *Foot and Ankle Orthopaedics*.2021May;6(2):24730114211013344. doi: 10.1177/24730114211013344.