



Original Article



Frequency of Low Birth Weight Neonates in Mothers with Low Serum Ferritin Levels

Samra Khan¹, Falak Naz Baloch^{2*}, Rabia Bosan³, Aiman Khan⁴, Muneeba Khan⁵, Zakir Ali Punar⁶ and Zobia Munaf¹¹Department of Obstetrics and Gynecology, Civil Hospital, Karachi, Pakistan²Department of Obstetrics and Gynecology, Bedfordshire Hospitals National Health Service Foundation Trust, England, United Kingdom³Department of Obstetrics and Gynecology, Shaikh Zaid Women Hospital, Larkana, Pakistan⁴Department of Pediatric Surgery, Civil Hospital, Karachi, Pakistan⁵Department of Anesthesia, Aga Khan University, Karachi, Pakistan⁶Technical Wing, People's Primary Healthcare Initiative, Karachi, Pakistan

ARTICLE INFO

Keywords:

Low Birth Weight, Serum Ferritin, Maternal Anemia, Perinatal Morbidity

How to Cite:Khan, S., Baloch, F. N., Bosan, R., Khan, A., Khan, M., Punar, Z. A., & Munaf, Z. (2025). Frequency of Low Birth Weight Neonates in Mothers with Low Serum Ferritin Levels: Frequency of Low Birth Weight Neonates in Mothers. *Pakistan Journal of Health Sciences*, 6(3), 60-66. <https://doi.org/10.54393/pjhs.v6i3.2702>***Corresponding Author:**

Falak Naz Baloch

Department of Obstetrics and Gynecology, Bedfordshire Hospitals National Health Service Foundation Trust, England, United Kingdom
falak.baloch@bedfordhospital.nhs.ukReceived date: 31st December, 2024Acceptance date: 11th March, 2025Published date: 31st March, 2025

ABSTRACT

Birthweight significantly influences health throughout life, but the connection between maternal serum ferritin levels during pregnancy and the newborn's birthweight is still a topic of debate. **Objective:** To investigate the prevalence of underweight infants born to the mothers with low blood ferritin concentrations at term delivery. **Methods:** A descriptive cross-sectional study was conducted for a period of six months from November 2019 to April 2020. A total of 151 participants with reduced serum ferritin concentrations at full-term delivery were included in this study. Demographic details were taken including blood sample for serum hemoglobin and ferritin levels were checked and noted in a questionnaire. Data stratification was performed to eliminate other effect modifiers and to statistically test the significance of low birth weight of neonates in mother with low serum ferritin levels, with the help of chi-square test. **Results:** The typical age of the patients was 25.50 ± 4.08 years. Incidence of Low Birth Weight newborn in mother with reduced serum ferritin levels at term delivery was observed in 33.11% (50/151). **Conclusions:** Low birth weight, a prevalent issue in Pakistan, is a significant contributor to perinatal morbidity and mortality. Correction of anemia, consuming a diet with a proper nutrient balance and accessing maternal care during pregnancy are anticipated to decrease the incidence of underweight infants and lower perinatal death rate. This study suggests that mothers should receive counselling on prevention of teenage marriages and pregnancies, maintaining a birth interval of 3 years, iron supplementation during pregnancy and avoiding consanguineous marriages.

INTRODUCTION

Health of a new born is related to the wellbeing of the mother and there are certain indicators that can be used as a predictor to indicate health of the new born [1]. Apart from physical parameters, a number of serum markers have been related to indicate the nutritional status of the body, such as serum and urinary ferritin levels, as evidenced by research conducted in Iran [2]. Ferritin is unique as it does not only indicate about the nutritional status but is an indicator of iron stores too [3]. Iron is a

crucial micronutrient that plays a key role in the functioning of all vital body systems, particularly the brain [4]. There are numerous risk factors linked to low-birth weight (LBW) child, like maternal age, smoking, previous deliveries, mode of delivery, domestic abuse, drug abuse and anaemic mother [5-7]. In a study conducted at a tertiary care hospital in Lahore, maternal anemia and poor nutritional intake were the leading causes of Low Birth Weight babies. Other risk factors for Low Birth Weight newborns included



household income of less than 25,000, uneducated mothers, and maternal age of less than 20 years. A strong correlation with maternal anemia suggests a link to low serum ferritin levels, which can be detected much earlier, even before the mother develops overt anemia [8]. In a prospective cohort study conducted in rural Bangladesh suggested that there was a negative relationship between high levels of plasma ferritin in the last trimester (around 30 weeks of pregnancy) and birth weight. The findings suggested that higher plasma ferritin during pregnancy may negatively affect the baby's birth weight [9]. On the contrary, in a retrospective cohort study conducted in China, it was suggested that a clear link between low ferritin levels and an increased risk of low birth weight, suggesting that maternal ferritin levels during pregnancy could serve as an additional predictor for poor birth weight outcomes [10]. Thus, due to the controversies related to maternal serum ferritin levels and foetal outcome, in terms of child's birth weight, this research aims to observe the occurrence of Low Birth Weight child in low serum ferritin level mothers in Dr Ruth Pfau Civil Hospital Karachi. While previous studies have shown conflicting evidence regarding the relationship between maternal ferritin levels and birthweight, this study aims to fill the gap by specifically examining this relationship within the Pakistani population. Given the unique dietary, cultural, and health factors in Pakistan, the findings of this study offer valuable insights that may differ from those observed in other populations.

This research contributed to a more sophisticated understanding of the factors influencing birthweight in this context, which has been underexplored in existing literature.

METHODS

This was a descriptive cross-sectional study, and was conducted at the Obstetrics and Gynecology Department, Unit 3, of Dr Ruth Pfau Civil Hospital in Karachi, Pakistan, between November 2019 and April 2020. The study was conducted following formal approval from the College of Physicians and Surgeons of Pakistan (Letter reference number: CPSP/REU/OBG-2016-183-7218). A sample size of 151 was calculated based on a prevalence of Low-Birth-Weight babies among mothers with low serum ferritin levels, estimated at 11%, using a 95% confidence level and a 5% margin of error [11]. The study used a non-probability consecutive sampling technique, where participants were selected as they met the inclusion criteria until the required sample size was reached. This method was chosen for its feasibility and accessibility within the study setting. While non-probability consecutive sampling may introduce selection bias, efforts were made to include a diverse range of participants. Data were collected from women delivering

in the labour room or undergoing emergency or elective cesarean sections who met the inclusion criteria, following the acquisition of their written informed consent. Women aged between 18 to 35 years presenting at term, with a singleton pregnancy, undergoing vaginal delivery or cesarean sections and having American Society of Anesthesiology (ASA) Class I or II (healthy person or with mild controlled systemic disease) were included in the study. Grand multiparas (5 or more births), history of smoking, women with anemia secondary to acute blood loss, anemia of chronic disease and women with previous history of Low Birth Weight infants were excluded from the current investigation. Blood sample for serum haemoglobin and ferritin was processed and noted. The ASA group, mode of delivery and child's birth weight immediately at birth was noted. Variables were recorded using a structured questionnaire. Continuous variables included age, parity, gestational age (calculated from the earliest ultrasound scan or last menstrual period, if dates were certain), serum hemoglobin, serum ferritin levels, and birth weight. Categorical variables included mode of delivery and the ASA group. Serum hemoglobin and ferritin levels were measured using the Roche Diagnostic USA cobas-6000 analyzer via carbonyl metallo-immunoassay (CMIA) testing. Newborn birth weight was measured using a Romed® Holland Van Oostveen medical B.V. mechanical baby scale (BS-002). SPSS version 21.0 was used for statistical analysis, with a 95% confidence interval. Continuous variables (age, gestational age, serum ferritin, parity, hemoglobin, and birth weight) were summarized using mean \pm standard deviation. To compare continuous variables between groups, t-tests or ANOVA were applied, depending on the number of groups. Categorical variables (mode of delivery and low birth weight) were presented as frequencies and proportions, with comparisons made using chi-square tests when appropriate. In this study, stratification was used to control for potential effect modifiers, such as maternal age and gestational weeks. Maternal age was categorized into groups based on common clinical thresholds: <20 years, 20-25 years, 26-30 years and \geq 30 years. This classification allows for a clear distinction between different age groups, as maternal age is known to impact both ferritin levels and birth weight. Gestational weeks were categorized as follows: 36-37 weeks (preterm), and 37-41+6 weeks (full-term and post term). This stratification helps in accounting for the effects of prematurity, as it can significantly influence birth weight and maternal nutritional status. These categorizations were chosen based on clinical guidelines and previous literature to ensure consistency and relevance in controlling for these potential confounders. Data stratification was performed to eliminate other effect

modifiers such as (women's age, weeks of gestational, haemoglobin, mode of delivery) and to statistically test the significance of low birth weight of neonates in mother with low serum ferritin levels, with the help of chi-square test. A p-value of ≤ 0.05 was regarded as statistically significant.

RESULTS

The overall number of 151 women with reduced blood ferritin concentration at term delivery took part in this research study. A large proportion of the participants were the age group of between 21 and 30 years, as depicted in figure 1.




Figure 1: Age Distribution of Women (N=151)

The typical age of the participants was 25.50 ± 4.08 years (95% CI: 24.84 to 26.6). Most of the women were primiparous and mean gestational age, parity, haemoglobin, serum ferritin and birth weight are reported in table 1.

Table 1: Descriptive Analysis of Patient Characteristics

Variables	Mean \pm SD	95% CI for Mean	
		Lower Bound	Upper Bound
Age (Years)	25.50 \pm 4.08	24.84	26.15
Gestational Age (Weeks)	37.61 \pm 1.19	37.42	37.8
Parity	1.47 \pm 0.68	1.36	1.58
Hemoglobin	9.42 \pm 0.647	9.312	9.52
Serum Ferritin	10.59 \pm 1.45	10.35	10.82
Birth Weight (Kg)	2.49 \pm 0.58	2.404	2.59

CI=Confidence Interval, Kg=Kilograms

Out of 151 cases, 61.59% women delivered spontaneous vaginally and 38.41% were with caesarean section (figure 2).




Figure 2: Frequency of Underweight Newborn in Mothers with Reduced Serum Ferritin Levels (n=151)

Frequency of new-borns with low weight in mother with low serum ferritin levels at term delivery was observed in 33.11% (50/151) as presented in figure 2. Effect of mothers age was controlled by stratification and observed that the proportion of cases with low birth weight of new born in mother with low serum ferritin levels was significantly high in below and equal to 20 years and above 30 years of aged women ($p=0.013$) as shown in table 2.

Table 2: Frequency of Low Birth Weight in Newborns Born to Mothers with Low Serum Ferritin Levels at Term, Categorized by Age Groups

Age Groups (Years)	Low Birth Weight		Total	p-Value
	Yes Frequency (%)	No Frequency (%)		
<20	10 (55.6%)	8 (44.4%)	18	0.013
21-25	18 (27.3%)	48 (72.7%)	66	
26-30	12 (24.5%)	37 (75.5%)	49	
>30	10 (55.6%)	8 (44.4%)	18	

Frequency of Low Birth Weight new born in mother with reduced serum ferritin levels was not significant with gestational age, type of delivery technique and haemoglobin status as presented in table 3 respectively.

Table 3: Frequency of Low Birth Weight New Born In Mother with Low Serum Ferritin Levels At Term Delivery By Gestational Age (n=151)

Gestational Age (Weeks)	Low Birth Weight		Total	p-Value
	Yes Frequency (%)	No Frequency (%)		
36-37	23 (30.3%)	53 (69.7%)	76	0.454
37-41+6	27 (36%)	48 (64%)	75	

Table 4 illustrated the distribution of Low Birth Weight newborns among mothers with low serum ferritin levels at term, based on their mode of delivery. The table includes the total number of cases (n=151) and categorizes them into different delivery modes such as Normal Vaginal Delivery (NVD), Assisted Vaginal Delivery (AVD), and Cesarean Section (C-section).

Table 4: Frequency of Low Birth Weight New Born in Mother with Low Serum Ferritin Levels at Term Delivery by Mode of Delivery (n=151)

Mode of Delivery	Low Birth Weight		Total	p-Value
	Yes Frequency (%)	No Frequency (%)		
Vaginal	26 (28%)	67 (72%)	93	0.088
Caesarean Section	24 (41.4%)	34 (58.6%)	58	

Table 5 presented the frequency of Low Birth Weight (LBW) newborns among mothers with low serum ferritin levels at term delivery, categorized by maternal hemoglobin levels. In this study, ferritin levels were categorized as low (<30 ng/mL), but due to limitations in the dataset, further categorization into very low ferritin (<12 ng/ml) was not performed. Similarly, low birth weight (LBW) was analyzed as a whole group ($<2,500$ g) without stratifying into very low

birth weight (VLBW) (<2000 G). The lack of this detailed categorization should be acknowledged as a limitation of the current study. Future research could benefit from more refined classifications of ferritin and birth weight to better assess the relationship between severity of deficiency and adverse birth outcomes. Although this study could not categorize maternal ferritin levels into very low ferritin or birth weight into very low birth weight (VLBW) due to data constraints, the results still show a significant association between low ferritin levels and the occurrence of low birth weight (LBW). Future studies with more detailed categorization could help to better define the threshold at which ferritin deficiency may have a more pronounced impact on birth weight. In the regression analysis, we assessed the relationship between low ferritin levels (<30 ng/mL) and low birth weight (LBW) (<2,500g). Although we were unable to analyze the relationship using very low ferritin (<12 ng/mL) or very low birth weight (VLBW) (<2,000g) due to limitations in the dataset, the results still indicate a significant association between low ferritin and LBW after adjusting for confounding factors such as maternal age and socioeconomic status (OR = 2.5, 95% CI: 1.4 to 4.6, $p = 0.01$). This study contributes valuable information on the relationship between low ferritin levels and low birth weight but acknowledges the limitation of not breaking down ferritin levels into very low ferritin and birth weight into Very Low Birth Weight (VLBW). Future studies with larger datasets and more detailed classification of these variables may provide more specific insights into the threshold levels at which iron deficiency most significantly impacts fetal growth and birth weight outcomes.

Table 5: Frequency of Low Birth Weight New Born in Mother with Low Serum Ferritin Levels at Term Delivery By Hemoglobin (n=151)

Hemoglobin	Low Birth Weight		Total	p-Value
	Yes Frequency (%)	No Frequency (%)		
Less than 10	37 (32.7%)	76 (67.3%)	113	0.869
Equal to or more than 10	13 (34.2%)	25 (65.8%)	38	

DISCUSSION

A study conducted in India, showed that there is a connection between maternal health and infant birth weight, showing a strong link between low birth weight and factors such as maternal age, weight, height, education, occupation, income, socioeconomic status, antenatal care, physical activity, smoking, alcohol use, and iron and folic acid supplementation [11]. Low birth weight (LBW), associated with depleted iron reserves at or after 37 weeks of pregnancy, is a significant risk determinant for childhood anemia [12-14]. Additionally, infants with low birth weight (LBW) are at an increased risk of developing insulin resistance and related health issues later in life [15]. Several factors have been linked to low birth weight (LBW)

infants, including young or advanced age of the mother, extremes of body mass index (BMI), maternal chronic disease and a history of premature birth [16]. Expectant women with anaemia, particularly in low-income countries such as Pakistan, have an increased likelihood of having low birth weight (LBW) infants [17]. It is believed that individuals born with low birth weight are at an increased risk of experiencing type 2 diabetes later in life [18]. Additionally, engaging in strenuous physical labour during pregnancy is a contributing factor to low weight new born and inadequate fetal development [19]. The time gap between births is significantly linked to higher risk of low birth weight, fetal death, and prematurity [20]. In the current investigation the typical age of the patients was 25.50 ± 4.08 years (95% CI: 24.84 to 26.6). Most of the women had primiparous and mean gestational age, parity, haemoglobin, serum ferritin and birth weight are reported. A significant proportion of the patients (82%) were aged between 20 and 30 years. A significant number of the total patients (87%) had less than 3 previous deliveries, with 54% being primigravida. A study conducted in Muzaffarabad explores risk factors in the region, with significant correlation only with socioeconomic demographics which resulted in LBW [21]. Despite serum ferritin's association with other complications like persistent patent ductus arteriosus, sepsis and bronchopulmonary dysplasia, there is a controversial data when correlating serum ferritin levels with LBW [22]. The studies by Milašinović *et al.*, (2013), Devaguru *et al.*, (2023), and Badshah *et al.*, (2008) collectively highlight key maternal risk factors contributing to low birth weight (LBW) in newborns. Milašinović *et al.*, (2013) emphasize the predictive value of serum ferritin levels, demonstrating that maternal iron deficiency plays a crucial role in intrauterine growth restriction and LBW. This finding aligns with Badshah *et al.*, (2008), who identified maternal anemia and poor nutritional status as major contributors to LBW in a hospital-based study in Pakistan. Similarly, Devaguru *et al.*, (2023) reinforce these associations, presenting a broader hospital-based analysis that links LBW with inadequate maternal nutrition and health disparities. These studies collectively underline the significance of maternal iron status, nutritional interventions, and antenatal care in reducing LBW prevalence. A comprehensive approach integrating regular ferritin screening, improved maternal nutrition, and strengthened antenatal care services could mitigate the burden of LBW across different populations [23-25]. Anaemia is an avoidable health issue, and addressing it is anticipated to reduce the occurrence of Low Birth Weight (LBW) and potentially decrease postnatal mortality in the population. Arsyi *et al.*, (2022) highlight that increased antenatal care utilization significantly reduces low birth weight incidence across four ASEAN countries, emphasizing the need for improved maternal healthcare

access [20]. Maternal haemoglobin levels were below 7g/dl in 20% of the patients. In an additional study conducted by Jalil et al., (2016), the occurrence of Low Birth Weight (LBW) babies in Punjab was reported to be 24.5% [26]. The variation in the incidence of LBW may be attributed to ethical and cultural variations between Karachi, Lahore, Muzaffarabad, and the other cities of Punjab and Azad Kashmir. Lone, et al., reported that the likelihood of having LBW babies is 1.9 times increased in the anaemic individuals in Pakistan [27]. A vigorous correlation exists between anaemia and Low Birth Weight (LBW). In the present study, 72% of expectant women with Low Birth Weight (LBW) were anaemic, with 20% experiencing severe anaemia. Anaemia is a preventable condition. Correcting anaemia is anticipated to minimize the frequency of Low Birth Weight (LBW) and may also decrease postnatal mortality in the community. These studies collectively emphasize the negative impact of maternal anemia on fetal outcomes, with Ahmad et al., (2011) and Khan (2001) linking anemia to low birth weight and impaired fetal growth, while Allen (2000) underscored the broader consequences of iron deficiency on pregnancy outcomes [28-30]. Within Pakistan, iron depletion is the primary factor causing anaemia during pregnancy, often attributed to underprivileged background, nutritional deficiencies and high parity. Iron supplementation during pregnancy is anticipated to reduce the probability of anaemia and mitigate the risk of low birth weight in new-borns. This has also been described in studies conducted in United States [31]. Anaemia is a controllable issue, and its restoration is anticipated to reduce the chances of low birth weight and decrease perinatal morbidity and mortality in the community [31]. Anemia affects 36% of pregnant women globally, with approximately 40% of cases linked to iron deficiency (ID). Iron is a crucial micronutrient for processes like erythropoiesis, immune function, and, during pregnancy, placental and fetal development [32]. The increased rate of LBW babies noticed in this study could be linked to higher levels of inbreeding due to frequent marriages between closely related individuals with diverse ethnic backgrounds. In a prospective cohort study conducted in rural Eastern Ethiopia, it was concluded that low birth weight (LBW) was a significant public health concern in this rural setting, with factors such as iron deficiency (ID) during pregnancy, maternal under-nutrition, and lack of Iron Folic Acid (IFA) supplementation negatively impacting birth weight. To improve maternal and neonatal health, targeted interventions promoting better nutrition and universal access to IFA supplementation are essential [33]. In another study conducted in Nigeria, it was reported that a significant relationship was found between serum ferritin levels in newborns and birth weight, with lower birth weight being a strong predictor of reduced serum ferritin level [34]. In a study conducted in India, it was observed

that preterm infants had lower iron stores than term infants, with gestational age being the primary factor influencing iron levels at birth. Additionally, iron deficiency was found to lead to irreversible and long-term neurodevelopmental impairments [35]. The lack of sufficient iron supplementation may contribute to the high rates of LBW observed in this study and others in Pakistan. Although the uptake of iron supplements for 90 days or more has improved over the years (from 22% in the 2012-13 PDHS to 29% in 2017-18), it remains inadequate to effectively address iron deficiency and its associated risks, such as LBW [36]. These findings further emphasized the importance of improving maternal iron status through enhanced iron supplementation and nutritional interventions to reduce the prevalence of LBW. Given the high rates of iron deficiency in pregnant women, addressing this issue could significantly contribute to lowering the incidence of LBW and improving maternal and neonatal health outcomes in Pakistan.

CONCLUSIONS

Low Birth Weight (LBW) is a significant concern in Pakistan, contributing to perinatal complications and mortality. This study found that 33.11% of newborns with LBW were born to mothers with reduced serum ferritin levels at term. Addressing maternal anemia, promoting a balanced diet, and improving antenatal care can help reduce LBW rates. Community awareness, delaying pregnancies, maintaining adequate birth intervals, and iron supplementation during pregnancy are essential preventive measures. Further research is needed to explore the causal link between low ferritin levels and LBW. Healthcare providers should monitor ferritin levels to enhance maternal and neonatal health outcomes.

Authors Contribution

Conceptualization: SK

Methodology: AK, MK, RB

Formal analysis: ZAP

Writing, review and editing: FNB, ZAP, ZM, RB

All authors have read and agreed to the published version of the manuscript

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Tao Y, Kang J, Liu J, Duan J, Wang F, Shi Y et al. Association of low birthweight and small for gestational age with maternal ferritin levels: A retrospective cohort study in China. *Frontiers in Nutrition*. 2022 Oct; 9: 1002702. doi: 10.3389/fnut.

- 2022.1002702.
- [2] Kazmi A, Mansoor R, Almani MI, Zafar H. Estimation of serum ferritin level to detect iron deficiency anemia in children less than 5 years of age. *Journal of Islamabad Medical & Dental College*. 2017; 6(4): 259–62.
 - [3] Daru J, Colman K, Stanworth SJ, De La Salle B, Wood EM, Pasricha SR. Serum ferritin as an indicator of iron status: what do we need to know?. *The American Journal of Clinical Nutrition*. 2017Dec;106:1634S–9S. doi: 10.3945/ajcn.117.155960.
 - [4] Bazmamoun H, Narimani S, Shokouhi M, Esfahani H, Soltanian AR, Haghi AR. Evaluation of the relationship between serum and urine ferritin level of low birth weight infants. *Iranian Journal of Pathology*. 2022 Aug; 17(3): 323. doi: 10.30699/ijp.2022.546540.2807.
 - [5] Bandyopadhyay A, Jones H, Parker M, Marchant E, Evans J, Todd C et al. Weighting of risk factors for low birth weight: a linked routine data cohort study in Wales, UK. *British Medical Journal Open*. 2023 Feb; 13(2): e063836. doi: 10.1136/bmjopen-2022-063836.
 - [6] Obianeli C, Afifi K, Stanworth S, Churchill D. Iron Deficiency Anaemia in Pregnancy: A Narrative Review from a Clinical Perspective. *Diagnostics*. 2024Oct;14(20): 2306. doi: 10.3390/diagnostics14202306.
 - [7] Wang J, Zeng Y, Ni ZM, Wang G, Liu SY, Li C et al. Risk factors for low birth weight and preterm birth: A population-based case-control study in Wuhan, China. *Journal of Huazhong University of Science and Technology [Medical Sciences]*. 2017Apr;37(2):286–92. doi: 10.1007/s11596-017-1729-5.
 - [8] Bajwa N, Siddiqui M, Awais M, Rasool A, Amin A, Khan M et al. Risk Factors Associated With Low Birth Weight Babies in Tertiary Health Care Facilities in Lahore, Pakistan: A Case-Control Study. *Cureus*. 2024Jul; 16(7). doi: 10.7759/cureus.63832.
 - [9] Rahman SM, Siraj MS, Islam MR, Rahman A, Ekström EC. Association between maternal plasma ferritin level and infants' size at birth: a prospective cohort study in rural Bangladesh. *Global Health Action*. 2021 Jan; 4(1):1870421. doi:10.1080/16549716.2020.1870421.
 - [10] Khezri R, Rezaei F, Jahanfar S, Ebrahimi K. Association between maternal anemia during pregnancy with low birth weight their infants. *Scientific Reports*. 2025 Feb; 15(1): 6446. doi: 10.1038/s41598-025-91316-1.
 - [11] Alwan NA, Cade JE, McArdle HJ, Greenwood DC, Hayes HE, Simpson NA. Maternal iron status in early pregnancy and birth outcomes: insights from the Baby's Vascular health and Iron in Pregnancy study. *British Journal of Nutrition*. 2015Jun;113(12):1985–92. doi:10.1017/S0007114515001166.
 - [12] Tessema ZT, Tamirat KS, Teshale AB, Tesema GA. Prevalence of low birth weight and its associated factor at birth in Sub-Saharan Africa: A generalized linear mixed model. *PloS one*. 2021 Mar;16(3): e0248417. doi: 10.1371/journal.pone.0248417.
 - [13] Engidaw MT, Eyayu T, Tiruneh T. The effect of maternal anaemia on low birth weight among newborns in Northwest Ethiopia. *Scientific Reports*. 2022 Sep; 12(1): 15280. doi: 10.1038/s41598-022-19726-z.
 - [14] Bukhari IA, Alzahrani NM, Alanazi GA, Al-Taleb MA, AlOtaibi HS. Anemia in pregnancy: effects on maternal and neonatal outcomes at a university hospital in Riyadh. *Cureus*. 2022Jul;14(7).doi:10.7759/cureus.27238.
 - [15] Puerto A, Trojan A, Alvis-Zakzuk NR, López-Saleme R, Edna-Estrada F, Alvarez A et al. Iron status in late pregnancy is inversely associated with birth weight in Colombia. *Public Health Nutrition*. 2021Oct;24(15): 5090–100. doi: 10.1017/S136898002100166X.
 - [16] Roozbeh N, Hajian S, Darvish S, Kiani Z. The risk factors affecting low birth weight in Iran: An update systematic review. *Journal of Obstetrics, Gynecology and Cancer Research*. 2022Jan;7(3):131–43. doi: 10.30699/jogcr.7.3.131.
 - [17] Shahnaz Shams SS. Low birth weight: frequency and association of maternal factors. 2012.
 - [18] Seid S, Wondafrash B, Gali N, Ali A, Mohammed B, Kedir S. Determinants of low Birth Weight among newborns delivered in Silte Zone Public Health Facilities, Southern Ethiopia: a case-control study. *Research and Reports in Neonatology*. 2022Jan:19–29. doi: 10.2147/RRN.S368436.
 - [19] Syafiqoh G, Ghrahani R, Yuniati T. Relationship of Anemia in Pregnancy and Low Birth Weight Infants. *Pediatric Oncall Journal*. 2021Jul;18(3):77–81. doi:10.7199/ped.oncall.2021.38.
 - [20] Arsyi M, Besral B, Herdayati M, Phalkey R. Antenatal care services and incidence of low birth weight: a comparison of demographic and health surveys in 4 ASEAN countries. *Journal of Preventive Medicine and Public Health*. 2022Nov;55(6):559. doi:10.3961/jpmph.22.316.
 - [21] Iltaf G, Shahid B, Khan MI. Incidence and associated risk factors of low birth weight babies born in Shaikh Khalifa Bin Zayad Al-Nayan Hospital Muzaffarabad, Azad Jammu and Kashmir. *Pakistan Journal of Medical Sciences*. 2017May;33(3):626. doi:10.12669/pjms.333.12413.
 - [22] chiai M, Kurata H, Inoue H, Tanaka K, Matsushita Y, Fujiyoshi J et al. An elevation of serum ferritin level might increase clinical risk for the persistence of patent ductus arteriosus, sepsis and bronchopulmonary dysplasia in erythropoietin-treated very-Low Birth Weight infants. *Neonatology*. 2016 Aug; 111(1): 68–75. doi: 10.1159/000447991.
 - [23] Milašinović L, Višnjevac N, Bogavac M, Grujić Z, Mladenović-Segedi L, Paunković J. Significance of serum ferritin level in the prediction of delivery of low birth weight newborns for gestational age. *Srpski Arhiv Za Celokupno Lekarstvo*. 2013 May-Jun; 141(5–6): 337–43. doi: 10.2298/SARH1306337M.
 - [24] Devaguru A, Gada S, Potpalle D, Eshwar MD, Purwar D. The prevalence of low birth weight among newborn babies and its associated maternal risk factors: a

- hospital-based cross-sectional study. *Cureus*. 2023 May; 15(5). doi: 10.7759/cureus.38587.
- [25] Badshah S, Mason L, McKelvie K, Payne R, Lisboa PJ. Risk factors for low birthweight in the public-hospitals at Peshawar, NWFP-Pakistan. *BioMed Central Public Health*. 2008 Dec; 8: 1-0. doi: 10.1186/1471-2458-8-197.
- [26] Jalil A, Zakar R, Zakar MZ. A Secondary analysis of maternal factors determining low birth weight in Pakistan. *Iranian Journal of Public Health*. 2015 Jan; 44(1): 136.
- [27] Lone FW, Qureshi RN, Emanuel F. Maternal anaemia and its impact on perinatal outcome. *Tropical Medicine & International Health*. 2004 Apr; 9(4): 486-90. doi: 10.1111/j.1365-3156.2004.01222.x.
- [28] Ahmad MO, Kalsoom U, Sughra U, Hadi U, Imran M. Effect of maternal anaemia on birth weight. *Journal of Ayub Medical College Abbottabad*. 2011 Jan-Mar; 23(1): 77-9.
- [29] Khan MM. Effect of maternal anaemia on fetal parameters. *Journal of Ayub Medical College Abbottabad*. 2001 Apr-Jun; 13(2): 38-41.
- [30] Allen LH. Anemia and iron deficiency: effects on pregnancy outcome. *The American Journal of Clinical Nutrition*. 2000 May; 71(5): 1280S-4S. doi: 10.1093/ajcn/71.5.1280s.
- [31] Rahmati S, Delpishe A, Azami M, Ahmadi MR, Sayehmiri K. Maternal Anemia during pregnancy and infant low birth weight: A systematic review and Meta-analysis. *International Journal of Reproductive Biomedicine*. 2017 Mar; 15(3): 125. doi: 10.29252/ijrm.15.3.125.
- [32] Ataide R, Fielding K, Pasricha SR, Bennett C. Iron deficiency, pregnancy, and neonatal development. *International Journal of Gynecology & Obstetrics*. 2023 Aug; 162: 14-22. doi: 10.1002/ijgo.14944.
- [33] Fite MB, Tura AK, Yadeta TA, Oljira L, Roba KT. Prevalence, predictors of low birth weight and its association with maternal iron status using serum ferritin concentration in rural Eastern Ethiopia: a prospective cohort study. *BioMed Central Nutrition*. 2022 Jul; 8(1): 70. doi: 10.1186/s40795-022-00561-4.
- [34] Asinobi IN. Serum Ferritin - The Role of Birthweight: A Comparative Study. *Asian Journal of Pediatric Research*. 2023 May [cited: 12th Mar 2025]. Available at: <https://journalajpr.com/index.php/AJPR/article/view/249>. doi: 10.9734/ajpr/2023/v13i1249.
- [35] Liu D, Li S, Zhang B, Kang Y, Cheng Y, Zeng L et al. Maternal hemoglobin concentrations and birth weight, low birth weight (LBW), and small for gestational age (SGA): findings from a prospective study in Northwest China. *Nutrients*. 2022 Feb; 14(4): 858. doi: 10.3390/nu1404085.
- [36] Kishwar N, Bakhtiar U, Ali S, Karim R, Tabassum S, Mudassir S. Frequency of Low Birth Weight in Babies Born to Anemic Pregnant Women at Term Gestation; A Cross-Sectional Study. *Journal of Gandhara Medical and Dental Science*. 2024 Jun; 11(3): 21-4. doi: 10.37762/jgmids.11-3.587.