



Original Article



Assessment of Periodontal Health in Patients with Alzheimer's disease in Karachi, Pakistan

Ujala Mushtaque¹, Muhammad Ibtahaj Khan¹, Rohan Tariq¹, Syeda Rijah¹, Hibba Batool Zaidi¹ and Abdullah Aslam¹¹Department of Periodontology, Jinnah Medical and Dental College, Karachi, Pakistan

ARTICLE INFO

Keywords:

Alzheimer's disease, Periodontal Health, Cognitive Impairment, Plaque and Gingival Index

How to Cite:Mushtaque, U., Khan, M. I., Tariq, R., Rijah, S., Zaidi, H. B., & Aslam, A. (2025). Assessment of Periodontal Health in Patients with Alzheimer's disease in Karachi, Pakistan: Periodontal Health in Alzheimer's Patients. *Pakistan Journal of Health Sciences*, 6(3), 23-27. <https://doi.org/10.54393/pjhs.v6i3.2709>***Corresponding Author:**Ujala Mushtaque
Department of Periodontology, Jinnah Medical and Dental College, Karachi, Pakistan
drujala@hotmail.comReceived date: 31st December, 2024Acceptance date: 10th March, 2025Published date: 31st March, 2025

ABSTRACT

Alzheimer's disease (AD), causes permanent loss of neurons and neural connections. **Objective:** To assess the periodontal health status of Alzheimer's disease (AD) patients by assessing plaque index, gingival condition, probing pocket depth, attachment levels, and bleeding on probing, and to correlate these findings with cognitive impairment levels as determined by Mini-Mental State Examination (MMSE) scores. **Methods:** A cross-sectional study, was conducted at Medicare Hospital between July and October 2024. Total 60 patients were recruited in the study. Hence plaque index, gingival condition, probing depth, attachment levels, and bleeding on probing were used to assess periodontal health, whereas MMSE scores were used to classify cognitive impairment. For statistical analysis, ANOVA and unpaired t-test analyses are carried out using SPSS version 24.0. All of the findings were deemed statistically significant if P was less than 0.05. **Results:** The study results revealed that the demographic and neurological parameters such as age, the distribution of genders, and the number of teeth of those patients suffering from Alzheimer's disease. Hence, MMSE scores were significantly lower, indicating substantial cognitive impairment ($P < 0.001$). As the severity of AD increases, there is a trend for all recorded variables to get worse ($p < 0.01$). PPD and CAL reveal more periodontal tissue loss in patients with severe AD, while PI and GI show more plaque accumulation and gingival inflammation. **Conclusions:** This study highlighted a strong link between worsening periodontal health and Alzheimer's severity, emphasizing the need for proactive dental care to improve patient well-being.

INTRODUCTION

The progressive neurodegenerative disease known as Alzheimer's disease in which there are irreversible neural connections. It causes loss of memory, decline in cognition, and ultimately significant functional reliance and mortality [1]. It is the most commonly occurring form of dementia which is usually found among elderly individuals approximately 60–80% globally [2]. Individuals in the advanced phases become reliant on caretakers, underscoring the significant financial and emotional strain on families and healthcare systems [3, 4]. Chronic inflammatory conditions, known as periodontal disease, affect the periodontium [5, 6]. This has arisen due to microbial dysbiosis in numerous researches [7]. Through alterations in the oral microbiota, which contains a wide variety of microorganisms in the oral cavity, Alzheimer's

disease is connected to poor oral health [8, 9]. Recent studies have demonstrated that patients suffering from dementia have more chances of getting Alzheimer's disease. In contrast, those who experienced dementia earlier in life had more severe forms of Alzheimer's disease [10, 11]. Oral pathogens such as *Treponema*, spirochetes, *Porphyromonas gingivalis*, and *Prevotella intermedia* are more prevalent in these patients [12]. According to one theory, these periodontal bacteria could enter the brain through various pathways such as blood-brain barrier, and cause direct damage [13, 14]. Hence, there is little information on how this association shows up in Asian communities, especially in places like Pakistan where oral and neurological health outcomes may be influenced by cultural, nutritional, and medical practices [15].



Furthermore, the frequency and severity of periodontal disease among Alzheimer's disease patients may be significantly influenced by socioeconomic characteristics, dental care accessibility, and knowledge of oral hygiene habits in Pakistani communities. By assessing periodontal health in Alzheimer's patients while taking into account regional sociocultural and healthcare aspects, this study fills a gap in the body of existing data. The results of this study may influence interdisciplinary techniques to better manage Alzheimer's disease, enhance oral healthcare procedures for neurodegenerative patients, and influence public health policy.

METHODS

The cross-sectional research was conducted at Medicare Hospital from July to October of 2024. ERB/JMDC/Approval# 00021124 was the reference number. Purposive sample was used to determine the periodontal health of Alzheimer's disease patients. Hence, clinical attachment loss (CAL) was used as the primary parameter as it is the most reliable predictor for the advancement of periodontal disease. Based on a 95% confidence interval and the anticipated prevalence of periodontal disease among AD patients, a total sample size of 60 participants was produced using Open Epi software. Participants who were 50 years of age or older and had a confirmed diagnosis of AD were eligible. Participants had to have at least 12 natural teeth to be recruited those with mental, behavioral, or systemic conditions that affect oral health assessment, those taking drugs that affect cognition, and those who had had periodontal therapy within the previous six months were not part of the study. The goal of the study was explained to each selected participant, informed consent was taken from them. Further, the participants were allowed to leave the study at any time. Various tests were of the participants was taken to evaluate their mental and health status. The diagnosis was further confirmed by MRI and CT scans. Mini-Mental State Examination (MMSE) was used to measure the cognitive impairment. Four groups were made of participants on the basis of severity of condition such as it was considered mild (score ≥ 25), moderate (scoring ≥ 11), severe (score ≤ 10), and normal (score ≥ 26). The periodontal health status of patients was assessed such as Index of plaque and gingivae, level of Clinical Attachment. Hence the severity of Periodontal Disease was determined by Probing Pocket Depth (PPD), and bleeding on probing (%BOP). The level of Oral hygiene was classified as good, fair, or poor on the basis of index of periodontal and gingiva. The thickness of dental plaque which is present on tooth surfaces is measured through plaque index, with scores ranging from 0 which means no plaque to 3 means excessive plaque. The redness, swelling, and bleeding upon

probing, with scores from 0 (representing healthy gingiva) to 3 (indicating severe inflammation) helps to examine the gingival inflammation through gingival index. Periodontal probe is used to measure the depth of periodontal pockets and patients with deeper pockets shows that they have advanced periodontitis. Clinical Attachment Level (CAL) measures periodontal tissue loss by determining the distance from the cemento-enamel junction to the base of the pocket. The percentage of sites exhibiting Bleeding on Probing (%BOP) reflects the severity of gingival inflammation, more chances of periodontal disease progression. To analyze the data, SPSS version 24.0 was used. The mean \pm SD was used to report descriptive statistics. The differences among the groups in the severity of AD were assessed using a one-way ANOVA, and the clinical and demographic characteristics of the two groups were compared using unpaired t-tests. A p-value of less than 0.05 was considered statistically significant.

RESULTS

The results of the study revealed demographic factors of participants suffering from Alzheimer's disease. Hence, MMSE scores were significantly lower, indicating substantial cognitive impairment ($P < 0.001$) as shown in table 1.

Table 1: Demographic Factors of Participants with Alzheimer's disease

Variables	Participants with Alzheimer's Disease AD	Participants who are Non-Cognitively Impaired	p-Value
Age (Mean \pm SD)	66.3 \pm 8.1	63.8 \pm 7.9	
Gender			
Male	27/60 = 45.0%	27/62 = 43.5%	
Female	33/60 = 55.0%	35/62 = 56.5%	
Total teeth present	15.5 \pm 3.9	16.7 \pm 4.1	
Status of Oral Hygiene			$<0.001^*$
Good	3/60 = 5.0%	10/62 = 16.1%	
Fair	18/60 = 30.0%	19/62 = 30.6%	
Poor	39/60 = 65.0%	33/62 = 53.2%	
Mini-Mental State Examination Scores	13.8 \pm 7.9	28.9 \pm 1.1	

* $P < 0.05$ is considered statistically significant

As shown in Table 2 the severity of AD increases, there is a trend for all recorded variables Plaque index, gingival index, Probing Pocket Depth, Clinical Attachment Level, and % Bleeding on Probing to get worse ($p < 0.01$). PPD and CAL reveal more periodontal tissue loss in patients with severe AD, while PI and GI show more plaque accumulation and gingival inflammation.

Table 2: Groupwise Assessments Periodontal Parameters

Assessment of Parameters	Participants with Alzheimer's Disease AD Normal (Mean \pm SD)	Participants with Alzheimer's Disease AD Mild (Mean \pm SD)	Participants with Alzheimer's Disease AD Moderate (Mean \pm SD)	Participants with Alzheimer's Disease AD Severe (Mean \pm SD)	p-Value
Plaque Index (PI)	1.42 \pm 0.30	1.95 \pm 0.22	2.60 \pm 0.18	3.52 \pm 0.35	< 0.01*
Gingival Index (GI)	0.60 \pm 0.20	1.12 \pm 0.25	1.68 \pm 0.22	2.30 \pm 0.30	< 0.01*
Probing Pocket Depth (PPD)	2.32 \pm 0.48	3.12 \pm 0.42	4.05 \pm 0.38	5.10 \pm 0.55	< 0.01*
Clinical Attachment Level (CAL)	2.72 \pm 0.52	3.52 \pm 0.38	4.50 \pm 0.45	5.63 \pm 0.57	< 0.01*
% Sites with Bleeding on Probing (%BOP)	31.00 \pm 5.70	39.10 \pm 5.30	57.20 \pm 7.00	69.50 \pm 12.70	< 0.01*

DISCUSSION

In recent times medical research has been greatly emphasizing the connection between periodontal health and Alzheimer's disease. According to the findings, Alzheimer's disease patients' periodontal health considerably declined as the illness worsened. These results demonstrate the complex connection between oral health and cognitive decline, which is in line with another research done worldwide [16]. One of the studies conducted in 2023 observed similar results [17]. Hence, due to their poor oral hygiene, the inflammatory markers were directly impacted as some of the variables help in the spread of severe periodontal parameters such as inflammation of gums and increased buildup of plaque [18]. The study conducted revealed a high correlation between the deterioration of periodontal diseases and the degree of cognitive impairment. Periodontal indicators showed clear distinctions across the cognitive phases (normal, mild, moderate, and severe Alzheimer's disease). Hence, similar results were observed in earlier research, in which results revealed that the severe Alzheimer's disease group's mean index of plaque and gingival were more than those of the normal group [19]. Patients with severe Alzheimer's disease frequently struggle to carry out everyday tasks, such as keeping their teeth clean. As these patients are unable to take sufficient nutritional intake, they cannot clean their mouths properly, and irregular brushing all contribute to the fast advancement of periodontal disease [20]. The results showed Probing Pocket Depth and Clinical Attachment Level were higher. This finding implies that periodontal tissue damage exacerbates with worsening cognitive deterioration. Similar results were revealed in which dementia patients had considerably greater PPD and CAL than non-demented controls. These alterations reflect the development of periodontal disease, in which microbial biofilm buildup and persistent inflammation progressively weaken the alveolar bone and connective tissues [21, 22]. Further, the results showed that the bleeding on probing also increased significantly in the severe stages. This represents the gingival inflammatory response which is a typical clinical sign of active periodontal disease. Gingival bleeding is a symptom of gingivitis, which is frequently the initial stage of periodontal disease, according to earlier research [23]. Gingival inflammation was more common which can

progress to more serious periodontal diseases and if left untreated, then bleeding on probing will be increased [24]. The main limitations of the study were the study design due to which the capacity to establish a causal relationship between the disease and periodontium was limited. Furthermore, the influence of additional variables that can shed more light on the variation in periodontal health among Alzheimer's patients, such as socioeconomic level, medication usage, and nutritional status, was not investigated. This study provides useful information for Pakistan, a country with less research on this topic. Further, the necessity of focused dental care interventions for those with Alzheimer's disease, was planned. Hence, counseling of caretakers regarding oral hygiene was essential to reduce the negative consequences of these patients related to poor periodontal health [25]. The results highlight how crucial it is to include dental care in Alzheimer's patients' treatment plans to improve oral health and maybe delay the rate of cognitive deterioration. Furthermore, studies are required to investigate therapeutic and preventative measures meant to enhance these people's dental and cognitive health outcomes.

CONCLUSIONS

The research underscores a strong link between deteriorating health of periodontium and the advancement of Alzheimer's disease. As cognitive function declines, the severity of periodontitis intensifies, driven by plaque accumulation, inflammation of the gums, and loss of tissue. It is crucial to emphasize the importance of oral healthcare, as inadequate periodontal health may hasten cognitive decline. Incorporating proactive dental care into treatment strategies and providing education to caregivers regarding oral hygiene can help in improving oral healthcare.

Authors Contribution

Conceptualization: UM, SR

Methodology: MIK

Formal analysis: UM, RT, HBZ

Writing, review and editing: HBZ, AA

All authors have read and agreed to the published version of the manuscript

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Kumar A, Sidhu J, Goyal A, Tsao JW, Doerr C. Alzheimer disease(nursing). 2025 Jan.
- [2] DeTure MA and Dickson DW. The neuropathological diagnosis of Alzheimer's disease. *Molecular Neurodegeneration*.2019Aug;14(1):32.doi:10.1186/s13024-019-0333-5.
- [3] Breijyeh Z and Karaman R. Comprehensive review on Alzheimer's disease: causes and treatment. *Molecules*.2020Dec;25(24):5789.doi:10.3390/molecules25245789.
- [4] Clemmensen TH, Busted LM, Søborg J, Bruun P. The family's experience and perception of phases and roles in the progression of dementia: An explorative, interview-based study. *Dementia*.2019 Feb; 18(2): 490-513. doi: 10.1177/1471301216682602.
- [5] Martínez-García M and Hernández-Lemus E. Periodontal inflammation and systemic diseases: an overview. *Frontiers in Physiology*.2021 Oct; 12: 709438. doi: 10.3389/fphys.2021.709438.
- [6] Kinane DF, Stathopoulou PG, Papapanou PN. Periodontal diseases. *Nature reviews Disease primers*. 2017 Jun; 3(1): 1-4. doi: 10.1038/nrdp.2017.38.
- [7] Kamer AR, Craig RG, Niederman R, Fortea J, de Leon MJ. Periodontal disease as a possible cause for Alzheimer's disease. *Periodontology 2000*. 2020 Jun; 83(1): 242-71. doi: 10.1111/prd.12327.
- [8] RG, Niederman R, Fortea J, de Leon MJ. Periodontal disease as a possible cause for Alzheimer's disease. *Periodontology 2000*.2020Jun83(1):242-71.doi: 10.1111/prd.12327.
- [9] Bui FQ, Almeida-da-Silva CL, Huynh B, Trinh A, Liu J, Woodward J et al. Association between periodontal pathogens and systemic disease. *Biomedical Journal*. 2019 Feb; 42(1): 27-35. doi: 10.1016/j.bj.2018.12.001.
- [10] Singh Rao SK, Harding A, Simmons T, Robinson S, Kesavalu L, Crean S. Oral inflammation, tooth loss, risk factors, and association with progression of Alzheimer's disease. *Journal of Alzheimer's disease*. 2014 Sep; 42(3): 723-37. doi: 10.3233/JAD-140387.
- [11] Sheppard O and Coleman M. Alzheimer's disease: etiology, neuropathology and pathogenesis. *Exon Publications*.2020 Dec:1-21.doi:10.36255/exonpublications.alzheimersdisease.2020.ch1.
- [12] Scheltens P, De Strooper B, Kivipelto M, Holstege H, Chételat G, Teunissen CE et al. Alzheimer's disease. *The Lancet*.2021Apr;397(10284):1577-90.doi:10.1016/S0140-6736(20)32205-4.
- [13] Dahlen G, Basic A, Bylund J. Importance of virulence factors for the persistence of oral bacteria in the inflamed gingival crevice and in the pathogenesis of periodontal disease. *Journal of Clinical Medicine*.2019 Aug; 8(9): 1339. doi: 10.3390/jcm8091339.
- [14] Harding A and Singh Rao SK. Periodontitis and dementia: a bidirectional relationship?.*Journal of Dental Research*.2022Mar;101(3):245-6.doi: 10.1177/00220345211043461.
- [15] Pisani F, Pisani V, Arcangeli F, Harding A, Singh Rao SK. The mechanistic pathways of periodontal pathogens entering the brain: The potential role of treponema denticola in tracing Alzheimer's disease pathology. *International Journal of Environmental Research and Public Health*.2022Jul;19(15):9386.doi:10.3390/ijerph19159386.
- [16] Beydoun MA, Beydoun HA, Hossain S, El-Hajj ZW, Weiss J, Zonderman AB. Clinical and bacterial markers of periodontitis and their association with incident all-cause and Alzheimer's disease dementia in a large national survey. *Journal of Alzheimer's disease*.2020 May; 75(1): 157-72. doi: 10.3233/JAD-200064.
- [17] Caserta MT, Caccioppo D, Lapin GD, Ragin A, Groothuis DR. Blood-brain barrier integrity in Alzheimer's disease patients and elderly control subjects. *The Journal of Neuropsychiatry and Clinical Neurosciences*.1998Feb;10(1):78-84.doi:10.1176/jnp.10.1.78.
- [18] Thomson WM and Barak Y. Tooth loss and dementia: a critical examination. *Journal of Dental Research*.2021 Mar; 100(3): 226-31. doi: 10.1177/0022034520957233.
- [19] Said-Sadier N, Sayegh B, Farah R, Abbas LA, Dweik R, Tang N et al. Association between periodontal disease and cognitive impairment in adults. *International Journal of Environmental Research and Public Health*. 2023 Mar; 20(6): 4707. doi: 10.3390/ijerph20064707.
- [20] Nadim R, Tang J, Dilmohamed A, Yuan S, Wu C, Bakre AT et al. Influence of periodontal disease on risk of dementia: a systematic literature review and a meta-analysis. *European journal of epidemiology*.2020Sep; 35: 821-33. doi: 10.1007/s10654-020-00648-x.
- [21] Ming Y, Hsu SW, Yen YY, Lan SJ. Association of oral health-related quality of life and Alzheimer disease: A systematic review.*The Journal of Prosthetic Dentistry*.2020Aug;124(2):168-75.doi:10.1016/j.prosdent.2019.08.015.
- [22] Alam MS, Maity MK, Nazmi AS, Alam MS, Ansari MS. Oral health issues and preventive measures in geriatric populations. *Journal of Pharmaceutical Negative Results*.2022Dec;13(10):2647-55.doi:10.47750/pnr.2022.13.S10.316.
- [23] Laugisch O, Johnen A, Buergin W, Eick S, Ehmke B, Duning T et al. Oral and periodontal health in patients with Alzheimer's disease and other forms of dementia-

a cross-sectional pilot study. *Oral Health & Preventive Dentistry*. 2021Apr;19:255-261.doi:10.3290/j.ohpd.b1248937.

- [24] Dioguardi M, Crincoli V, Laino L, Alovise M, Sovereto D, Mastrangelo F *et al.* The role of periodontitis and periodontal bacteria in the onset and progression of Alzheimer's disease: a systematic review. *Journal of Clinical Medicine*. 2020Feb;9(2):495.doi:10.3390/jcm9020495.
- [25] de Oliveira Araújo R, Villoria GE, Luiz RR, Esteves JC, Leão AT, Feres-Filho EJ. Association between periodontitis and Alzheimer's disease and its impact on the self-perceived oral health status: a case-control study. *Clinical Oral Investigations*. 2021Feb;25:555-62. doi: 10.1007/s00784-020-03489-w.
- [26] Marchini L, Ettinger R, Caprio T, Jucan A. Oral health care for patients with Alzheimer's disease: An update. *Special Care in Dentistry*. 2019 May; 39(3):262-73.doi: 10.1111/scd.12375.