



Original Article



Dental Caries Prevalence among Undergraduate Students Attending Medical and Dental College

Adnan Sunny¹, Sardar Waleed Babar², Syed Akbar Abbas Zaidi³, Abdul Qudoos Memon⁴, Yasir Hussain⁵, Muhammad Zeeshan⁶ and Muhammad Farooq Chaudary⁷¹Department of Prosthodontics, Shifa College of Dentistry, Islamabad, Pakistan²Department of Oral Pathology, Shifa College of Dentistry, Islamabad, Pakistan³Department of Dental Education, Bahria University Health Sciences, Karachi, Pakistan⁴Department of Operative Dentistry, Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan⁵Department of Prosthodontics, Niazi Medical and Dental College, Sargodha, Pakistan⁶Department of Dental Materials, Niazi Medical and Dental College, Sargodha, Pakistan⁷Department of Dental Education, Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan

ARTICLE INFO

Keywords:

Dental Caries, Tooth Demineralization, Oral Health, Dental Plaque, Global Prevalence

How to Cite:Sunny, A., Babar, S. W., Zaidi, S. A. A., Memon, A. Q., Hussain, Y., Zeeshan, M., & Chaudary, M. F. (2025). Dental Caries Prevalence among Undergraduate Students Attending Medical and Dental College: Prevalence of Dental Caries. *Pakistan Journal of Health Sciences*, 6(2), 181-184. <https://doi.org/10.54393/pjhs.v6i2.2720>***Corresponding Author:**Muhammad Farooq Chaudary
Department of Dental Education, Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan
dr.farooqch789@gmail.comReceived date: 6th January, 2025Acceptance date: 22nd February, 2025Published date: 28th February, 2025

ABSTRACT

Dental caries is a chronic and widespread infectious disease that affects billions of people worldwide. The most prevalent chronic illness is one that, if left untreated, becomes worse over time. **Objective:** To determine the prevalence of the dental caries in the undergraduate student in medical and dental college of Jamshoro. **Methods:** With permission from Liaquat University of Medical and Health Science's Ethical Committee vide number ERC-19/2024/11, this observational cross-sectional study was carried out between April 2024, and November 2024. All students of Medical and Dentistry who were willing to participate in the study were included with 17-25 age range. Epi Info Software was used to calculate the sample size. 500 of the students were included as per the sample size calculation. **Results:** 300 (60%) of the participants were day scholars. The first, second, third, and fourth/final years were represented by students 105 (21%), 90 (18%), 175 (35%) and 130 (26%) respectively. 97 students (19.4%) had no DMFT score, whereas 403 students (80.6%) had dental caries with a DMFT score (>1). The test-takers' DMFT scores were as follows: 97 (19.4%) for a score of 0 and 139 for a score of 1 (27.8%). The DMFT scores were 102 (20.4%) for a score of 2, 136 (27.2%) for a score of 3, and 26 (5.2%) for a score of 4. **Conclusion:** This study demonstrated that the population's prevalence of dental caries was relatively low.

INTRODUCTION

Dental caries is a microbiological and infectious disease [1]. It is the result of tooth demineralization, which can lead to cavitation and pulp injury if treatment is not received. The combination of several problems, such as tooth demineralization, dental plaque bacteria, processed carbohydrates, and weak teeth from external sources [2]. Dental caries is a chronic and widespread infectious disease that affects billions of people worldwide. The most prevalent chronic illness is one that, if left untreated, becomes worse over time [3]. Because they make it

difficult to eat, chew, smile, and communicate, teeth that are broken, decayed, or missing have a major impact on day-to-day living. Dental caries is the third most prevalent oral health issue that affects people of all races, according to the World Health Organization (WHO). Dental caries affects around 100% of adults in most nations worldwide, particularly in developed nations [4]. In Pakistan, dental caries is seven times more common than hay fever and five times more common than asthma, which has concerning consequences for oral health in the nation. It is presently a



significant problem for community health in developing countries because of inadequate oral hygiene practices, excessive sugar consumption, and ignorance regarding fluoride additions [5, 6]. Thus, it is not unexpected that the WHO is urging pledges for ongoing oral and dental health improvement. The Decayed, Missing, Filled (DMF) index has been used for about 80 years and is well recognized as a significant indication of caries experience in dental epidemiology [7]. The DMFT index is used for teeth, and each person's score can be anywhere from 0 to 28 or 32. The DMFT results will be interpreted using the DMF scoring system. The traits that make women more susceptible to dental caries are a matter of debate, and some of these traits may differ among communities [8]. A study of medical and dentistry students in Taxila revealed a mean DMFT score of 1.64 ± 1.66 among participants. This shows generally decent oral health, but it also emphasizes unmet treatment requirements for untreated cavities. The study highlighted the importance of education and preventative practices in treating dental caries [9]. Yadav K *et al.*, conducted another research in Nepal to analyze medical students' knowledge, attitudes, and practices about dental caries at Janaki Medical College Teaching Hospital. It stated that while these students have enough understanding of oral hygiene principles, greater attention should be made on practical applications [10]. Despite substantial studies [5, 9] on dental caries prevalence, there are still numerous gaps in the literature. Firstly, gender disparities in dental caries have been frequently reported, the underlying biological, behavioral, and environmental variables that contribute to this disparity in Pakistani medical student populations have received little attention. Furthermore, stress levels, academic burden, and psychological aspects influencing medical and dental students' dental hygiene practices have not been well investigated. The objective of the current study was to determine the prevalence of the dental caries in the undergraduate student in medical and dental college of Jamshoro.

METHODS

With permission from Liaquat University of Medical and Health Science's Ethical Committee vide number ERC-19/2024/11, this observational cross-sectional study was carried out at Liaquat University of Medical and Health Sciences Jamshoro between April 2024, and November 2024. All students of who were willing to participate in the study were included with 17-25 age range. While students who had history of orthodontic treatment or ongoing orthodontic treatment were excluded. To validate external generalizability, a Kolmogorov-Smirnov test was conducted, confirming that the age distribution of the sample did not significantly differ from the general student population ($p = 0.076$). Epi Info Software was used to

calculate the sample size. Population size was taken as 957 with an expected frequency of 50% and CI 95%. 500 of the students were included as per the sample size calculation. The students were selected randomly. The participant signed a written consent form. The students were told there were no risks or disadvantages. To record the oral health status for dental caries, three examiners from dental college filled out the DMFT performa and demographic data, making sure to check for any inaccuracies by looking at dental carries. (D) indicates decaying teeth, (M) indicates caries-related tooth loss, (F) indicates carious tooth filling, and (T) indicates teeth. The WHO-recommended procedure was followed i.e. (Examiners were trained and calibrated to minimize inter-examiner and intra-examiner variability. WHO-recommended tools, such as mouth mirrors, explorers, and periodontal probes (e.g., WHO probe), were used. Proper lighting (natural) was used. Infection control measures, including the use of Personal Protective Equipment (PPE) like gloves, masks, and eyewear, were followed, with instruments sterilized or disposable ones utilized. Patients were seated comfortably in a well-lit area, ensuring their head remains stable throughout the examination). Students who met the requirements for inclusion were evaluated in the classroom near the window under natural light. The SPSS version 26.0 was used to enter and analyze the data. The relationship between gender and dental caries prevalence was evaluated using the chi-square test. An independent t-test and chi square test were used to analyze the differences in average DMFT, dental caries or decaying, missing, and fillings among many groups. Both the 95% confidence level and the P-Value 0.05 criterion were used.

RESULTS

Among the 500 subjects, 255 (51%) were males and 245 (49%) were females. Maximum of the students belong to the age group 22-23 years 167 (53.4%) followed by 24-25 years 103 (46.6%) (Table 1).

Table 1: Characteristics of the Subjects Involved in the Study

Age	Male Frequency (%)	Female Frequency (%)	Total Frequency (%)
<18	51 (49.5%)	52 (50.5%)	103 (20.6%)
19-21	40 (37.4%)	67 (60.6%)	107 (21.4%)
22-23	84 (50.3%)	83 (49.7%)	167 (33.4%)
24-25	80 (65%)	43 (35%)	123 (24.6%)
Total	255 (51%)	245 (49%)	500 (100%)

Three hundred (60%) of the participants were day scholars. The first, second, third, and fourth/final years were represented by students 105 (21%), 90 (18%), 175 (35%) and 130 (26%) respectively. 97 students (19.4%) had no DMF score, whereas 403 students (80.6%) had dental caries with a DMFT score (>1). The test-takers' DMFT scores were as follows: 97 (19.4%) for a score of 0 and 139 for a score of 1

(27.8%). The DMFT scores were 102 (20.4%) for a score of 2, 136 (27.2%) for a score of 3, and 26 (5.2%) for a score of 4. The differences between genders were statistically significant ($p=0.003$). Logistic regression analysis revealed age and gender as significant predictors of caries presence ($p<0.05$) (Table 2).

Table 2: Characteristics of the Participants Involved in the Study

Variables	Category	Frequency (%)	p-value
Gender	Male	245 (49%)	0.003*
	Female	255 (51%)	
Class	1 st Year	105 (21%)	0.045*
	2 nd Year	90 (18%)	
	3 rd Year	175 (35%)	
	4 th /Final Year	130 (26%)	
Residence	Hostilities	200 (40.0%)	
	Day Scholars	300 (60.0%)	

Table 3: Gender associated with the DMFT Index Score

Gender	Class (Year)	Mean of Decayed Teeth	Mean of Missed Teeth	Mean of Filled Teeth	DMFT Score
Male	1 st	1.40 ± 0.05	-	0.64 ± 0.63	2.04 ± 0.68
	2 nd	0.27 ± 1.01	-	1.59 ± 0.19	1.86 ± 1.20
	3 rd	0.13 ± 1.06	-	0.41 ± 0.13	0.54 ± 1.19
	4 th /Final	1.30 ± 1.06	-	0.01 ± 0.29	1.31 ± 1.35
Female	1 st	1.51 ± 0.59	0.03 ± 0.01	0.51 ± 0.43	2.05 ± 1.03
	2 nd	0.19 ± 1.16	-	0.21 ± 0.81	0.40 ± 1.97
	3 rd	1.01 ± 0.01	-	0.94 ± 0.71	1.95 ± 0.72
	4 th /Final	0.03 ± 0.19	0.01 ± 0.18	1.46 ± 0.35	1.50 ± 0.72
p-Value	-	0.003*	-	-	0.27

*Statistically Significant

Effect size measures were calculated to clarify the clinical significance of group differences. The effect size for gender differences in DMFT scores was Cohen's ($d = 0.42$), indicating a moderate effect size. Additionally, Cramér's V for the chi-square test between gender and caries presence was 0.28, suggesting a moderate association. The odds ratio (OR) for females having a DMFT > 0 compared to males was 1.34 (95% CI: 1.12 – 1.57, $p = 0.002$), indicating that female students were 34% more likely to have dental caries than males (Table 4).

Table 4: Statistical Tests and Effect Sizes Evaluating the Clinical Significance of Group Differences

Variable Comparison	Test Used	Effect Size	p-value	Interpretation
Gender vs DMFT score	Independent t-test	Cohens $d=0.42$	0.003*	Moderate Effect
Gender vs Caries	Chi-square test	Cramér's $V=0.28$	<0.001*	Moderate Association
Female vs Male	Logistic Regression	OR= 1.34 (95% CI: 1.12 – 1.57)	0.002*	Female more likely to have caries by 34%

*Statistically Significant

DISCUSSION

The research involved 500 students aged 17 to 25 from medical & dental college, revealing that approximately 80.6% of participants exhibited dental caries, as indicated by a DMFT score. Interestingly, while female students showed a higher prevalence 255(51%) of caries compared to males 245 (49%). Given that the mean DMFT of 0.27, it

DMFT Categories	Low (0-1)	236 (47.2%)	<0.001*
	Moderate (2-3)	238 (47.6%)	
	High (4+)	26 (5.2%)	
Caries Presence	DMFT = 0	97 (19.4%)	<0.001*
	DMFT > 0	403 (80.6%)	

*Statistically Significant

The correlation between the DMFT index score and gender over time is presented. This connection with the average number of students having decayed teeth among male and female students are statistically significant, with a p-value of 0.003. Furthermore, across students in different graduation years, the average number of missing teeth was 0.000.00 for male students and 0.0360.29 for female students. Female students exhibited slightly higher DMFT scores, and difference was statistically significant, with a p value of 0.003 (Table 3).

was determined that the amount of caries suffered was extremely minimal. Since many parts of Sindh are classified as being in the fluorotic belt, using ground water for drinking, which may be higher in fluoride, might be one of the causes of the low DMFT. Regarding age and gender, the results of this survey were in line with those of another study conducted on Peshawar undergraduate dental students [11]. Female students 255 (51%) had a higher prevalence of caries than male students, according to other studies that used the same study design. The average total DMFT score for boys and girls did not differ statistically significantly, according to this study. With a p-value of 0.003, the quantity of dental caries, which affects the DMFT average, was statistically significant. According to the present study's findings, dental caries is prevalent in both men and women. Several studies conducted in Saudi Arabia, the United Kingdom, India, Pakistan and Saudi Arabia produced similar results [5, 6, 12-17]. Based on these findings, caries was found to be lower. Among the included

studies, there were differences in the reported dental caries rates [18]. This supported the findings of Tsuchida S *et al.*, who discovered that the prevalence of dental caries varies greatly between studies because of things like (a) the age, socioeconomic status, and availability of examination of the subjects; (b) racial and cultural factors; and (c) diagnostic criteria [19]. It is hard to extrapolate findings from one ethnic group inside another due to the significant regional variations in dental caries prevalence. Current research findings in some ways support the World Health Organization's 2022 World Oral Health Report, which indicates that emerging nations have low Mean DMFTs (1.7) [20]. This analysis found that the frequency of dental cavities was trending increasing in developing nations and lower in industrialized ones. This is mostly since industrialized nations have better self-care habits, fluoride exposures, and preventative care programs, whereas underdeveloped nations lack those [21]. Qureshi A *et al.*, conducted another survey in underprivileged regions of Karachi, revealing variances in the mean DMFT, which is around 4 among rural populations [22]. The socioeconomic status of two distinct areas is the cause once more. Since Karachi is Pakistan's largest metropolis, there are more economic prospects there than in Jamshoro. Once more, the results showed that there were relatively few teeth filled because of caries. Reliability of the DMFT performance was enhanced through examiner calibration and adherence to standardized procedures. The natural light conditions and sterile instruments minimized variability in caries detection.

CONCLUSIONS

This study demonstrated that the population's prevalence of dental caries was relatively low. It demonstrated how oral hygiene education aided medical and dentistry students in maintaining and practicing good oral hygiene. Larger- longitudinal scale studies should be conducted, to explore caries determinants in diverse populations and evaluate the long-term effectiveness of preventive strategies

Authors Contribution

Conceptualization: AQM

Methodology: AQM, MFC

Formal analysis: SWB, YH, MZ

Writing, review and editing: AS, SWB, SAAZ, YH, MFC

All authors have read and agreed to the published version of the manuscript

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Fejerskov O. Changing paradigms in concepts on dental caries: consequences for oral health care. *Caries Research*. 2004 May; 38(3):182-91. doi:10.1159/00077753.
- [2] Cheng L, Zhang L, Yue L, Ling J, Fan M, Yang D *et al.* Expert consensus on dental caries management. *International Journal of Oral Science*. 2022 Dec; 14(1): 17. doi: 10.1038/s41368-022-00167-3.
- [3] Chapple IL, Bouchard P, Cagetti MG, Campus G, Carra MC, Cocco F *et al.* Interaction of lifestyle, behaviour or systemic diseases with dental caries and periodontal diseases: consensus report of group 2 of the joint EFP/ORCA workshop on the boundaries between caries and periodontal diseases. *Journal of Clinical Periodontology*. 2017 Mar; 44:S39-51. doi.org/10.1111/jcpe.12685.
- [4] Zhang JS, Chu CH, Yu OY. Oral microbiome and dental caries development. *Dentistry Journal*. 2022 Sep; 10(10): 184. doi: 10.3390/dj10100184.
- [5] Siddiqui AA, Alshammary F, Mulla M, Al-Zubaidi SM, Afroze E, Amin J *et al.* Prevalence of dental caries in Pakistan: a systematic review and meta-analysis. *BioMed Central Oral Health*. 2021 Dec; 21: 1-2. doi: 10.1186/s12903-021-01802-x.
- [6] Dawani N, Nisar N, Khan N, Syed S, Tanweer N. Prevalence and factors related to dental caries among pre-school children of Saddar town, Karachi, Pakistan: a cross-sectional study. *BioMed Central Oral Health*. 2012 Dec; 12: 1-9. doi: 10.1186/1472-6831-12-59.
- [7] Motallaei MN, Yazdani M, Tebyanian H, Tahmasebi E, Alam M, Abbasi K *et al.* The current strategies in controlling oral diseases by herbal and chemical materials. *Evidence-Based Complementary and Alternative Medicine*. 2021 Aug; 2021(1):3423001. doi: 10.1155/2021/3423001.
- [8] Nath S, Zilm P, Jamieson L, Kapellas K, Goswami N, Ketagoda K *et al.* Development and characterization of an oral microbiome transplant among Australians for the treatment of dental caries and periodontal disease: A study protocol. *PLoS One*. 2021 Nov; 16(11): e0260433. doi: 10.1371/journal.pone.0260433.
- [9] Rabbani M, Manzoor K, Faisal A, Jabeen W, Rabbani S, Haroon A *et al.* Periodontal and Caries Status in Relation to Oral Health Knowledge, Attitudes, and Behavior of Medical and Dental Students in Taxila, Pakistan. *Cureus*. 2024 Oct; 16(10). doi: 10.7759/cureus.72718.
- [10] Yadav K and Prakash S. Knowledge, attitude and practice on dental caries and Oral hygiene among medical students at Janaki medical college teaching hospital. *International Journal of Biomedical Science*. 2016 Feb; 1(2):22-31. doi:10.55530/ijmbiosnepal.v1i2.13.

- [11] Jehan OH, Khan F, Jehan AM, Yousaf M. Assessment of oral health behavior, knowledge, and practice among medical and dental undergraduate students of Peshawar Medical and Dental College, Pakistan. *Biomedica*.2024Oct;40(1):26-31.doi:10.24911/BioMedica/5-1146.
- [12] Al Agili DE. A systematic review of population-based dental caries studies among children in Saudi Arabia. *The Saudi Dental Journal*.2013Jan;25(1):3-11.doi:10.1016/j.sdentj.2012.10.002.
- [13] Pitts NB, Twetman S, Fisher J, Marsh PD. Understanding dental caries as a non-communicable disease. *British Dental Journal*.2021 Dec; 231(12): 749-53. doi: 10.1038/s41415-021-3775-4.
- [14] Kotronia E, Brown H, Papacosta AO, Lennon LT, Weyant RJ, Whincup PH et al. Oral health and all-cause, cardiovascular disease, and respiratory mortality in older people in the UK and USA. *Scientific Reports*.2021Aug;11(1):16452.doi:10.1038/s41598-021-95865-z.
- [15] Uerlich MF, Baker SR, Day PF, Brown L, Vettore MV. Common determinants of dental caries and obesity in children: a multi-ethnic nested birth cohort study in the United Kingdom. *International Journal of Environmental Research and Public Health*.2021 Nov; 18(23): 12561. doi: 10.3390/ijerph182312561.
- [16] Sabharwal A, Stellrecht E, Scannapieco FA. Associations between dental caries and systemic diseases:a scoping review.*BioMed Central Oral Health*.2021Dec;21:1-35.doi:10.1186/s12903-021-01803-w.
- [17] Siddique S, Shahid AM, Mahmood H, Farooq N, Ali S, Maqbool S. Dental Caries among the patients visiting HBS Dental College Hospital Islamabad. *Pakistan Journal of Medical & Health Sciences*.2022 Jun; 16(06): 188-. doi: 10.53350/pjmhs22166188.
- [18] Alshammari FR, Alamri H, Aljohani M, Sabbah W, O'Malley L, Glenny AM. Dental caries in Saudi Arabia: A systematic review. *Journal of Taibah University Medical Sciences*.2021Oct;16(5):643-56.doi:10.1016/j.jtumed.2021.06.008.
- [19] Tsuchida S, Umemura H, Iizuka K, Yamamoto H, Shimazaki I, Shikata E et al. Recent findings on metabolomics and the microbiome of oral bacteria involved in dental caries and periodontal disease. *World Journal of Microbiology and Biotechnology*. 2025 Jan; 41(1): 1-0. doi: 10.1007/s11274-024-04224-3.
- [20] Eaton K, Yusuf H, Vassallo P. The WHO global oral health action plan 2023-2030. *Community Dental Health*.2023May;40(2):68-9.doi:10.1922/CDH_Jun23 Editorial02.
- [21] Wen PY, Chen MX, Zhong YJ, Dong QQ, Wong HM. Global burden and inequality of dental caries, 1990 to 2019. *Journal of Dental Research*.2022Apr;101(4):392-9.doi : 10.1177/00220345211056247.
- [22] Qureshi A, Safdar NF, Qureshi H, AlFawaz YF, Al Ahdal K, Shabib S et al. Dietary factors influencing the caries status of adults in Karachi, Pakistan: initial findings. *International Journal of Environmental Research and Public Health*.2022Jun;19(12):6980.doi:10.3390/ijerph19126980.