



Original Article



Beyond Weight Loss: Investigating Irritability and Mood Swings after Bariatric Surgery: A Prospective Study

Mushtaq Ahmad¹, Maria Alamgir¹, Waqar Ahmad¹, Mah Noor Dad Khan¹ and Abdul Haseeb Sahibzada¹¹International Metabolic and Bariatric Center, Afridi Medical Complex, Peshawar, Pakistan

ARTICLE INFO

Keywords:

Bariatric, Surgery, Irritability, Postoperative Complications

How to Cite:Ahmad, M., Alamgir, M., Ahmad, W., Khan, M. N. D., & Sahibzada, A. H. (2025). Beyond Weight Loss: Investigating Irritability and Mood Swings after Bariatric Surgery: A Prospective Study: Weight Loss: Investigating Irritability and Mood Swings After Bariatric Surgery. *Pakistan Journal of Health Sciences*, 6(6), 42-48. <https://doi.org/10.54393/pjhs.v6i6.3007>***Corresponding Author:**Waqar Ahmad
International Metabolic and Bariatric Center, Afridi Medical Complex, Peshawar, Pakistan
dr.waqar84@gmail.comReceived Date: 25th March, 2025Revised Date: 6th June, 2025Acceptance Date: 11th June, 2025Published Date: 30th June, 2025

ABSTRACT

The physical changes after bariatric surgery are evident, but the significant changes in psychological and emotional effects are underscored and have not received much attention. **Objectives:** To determine the prevalence of Irritability and Mood Swings After Bariatric Surgery and its impact on hormonal and nutritional changes. **Methods:** This prospective observational study was conducted from February 2022 to January 2023. Patients undergoing any type of bariatric surgery with no existing psychiatric conditions that required medication were included. Structured questionnaires were distributed among the participants, and informed consent was also provided. **Results:** The study comprised 1,014 patients, with a mean age of 33 ± 13 years. According to the Profile of Mood States (POMS), 679 (67%) of participants had clinically severe mood abnormalities before surgery, which then increased to 844 (83%) at three months and then reduced to 355 (35%) at six months ($p < 0.001$). Compared to 740 (73%) before surgery, 902 (89%) of patients experienced moderate to severe irritation at three months ($p < 0.001$). After six months, this percentage dropped to 446 (44%). **Conclusions:** The research concluded notable changes in mood, irritability, and hormone levels after bariatric surgery, including increased irritability and mood disturbance at three months, with improvement at six months. Prevalent nutritional deficiencies of vitamins B12 and D regression were also found to be associated with increased mood and irritability. Multiple hormonal variations and nutritional deficiencies were found to result in postoperative mood variation significantly.

INTRODUCTION

Bariatric surgery has become a revolutionary treatment option and is considered a very viable alternative that can lead to substantial and long-lasting weight loss for those suffering from obesity [1]. This surgical strategy includes several operations that change the gastrointestinal tract's structure, which lowers food consumption and nutritional absorption [2]. Although the physical changes post-bariatric surgery is usually the main outcome of interest, the psychological and emotional changes following surgery are just as significant and yet are often overlooked, and research on the psychological effects of bariatric surgery has not received much attention [3]. Among the many psychological aspects of bariatric surgery, irritability and mood swings stand out as prevalent but little-known issues

that can have a big influence on a patient's long-term success [4]. Following surgery, certain patients exhibit a higher chance of experiencing negative mental health consequences [5]. High rates of preexisting depression, unfulfilled expectations about the transformative power of weight loss, and post-surgical changes in the hormonal and nutritional levels are some of the likely contributing factors leading to these adverse mental health outcomes, including mood changes and irritability [6]. After bariatric surgery, there are a variety of intricately linked physiological and psychological changes [7]. A key factor is the hormonal changes brought on by abrupt weight loss and changes in the gastrointestinal tract. RYBG and SG result in the enhancement of postprandial secretion of



satiety hormones, including glucagon-like peptide-1 (GLP-1) and oxyntomodulin (OXM), which contribute to a feeling of fullness and decrease hunger, resulting in the postoperative appetite regulation [8]. Following bariatric surgery, hormones including ghrelin, leptin, insulin Peptide YY (PYY) play a vital role in maintaining weight loss [9]. Ghrelin, a hunger hormone, also decreased after surgery due to the removal of ghrelin-secreting gastric tissue [10]. Leptin, produced by adipose tissue, sends feedback signals to the hypothalamus and regulates regular appetite and balances energy [11, 12]. The enteroendocrine changes brought on by sleeve gastrectomy and Roux-en-Y gastric bypass (RYGB) have an impact on mental health, owing to the evidence now available, which indicates that several hormones and neuropeptides that regulate hunger have an impact on mood [13]. Similarly, a deficiency of vital minerals such as iron, magnesium, vitamin B12, and vitamin D can impair neurological function and lead to mood swings [14, 15]. Patients frequently struggle with adjusting to new eating habits and managing their altered body image, which can cause them to feel frustrated, anxious, or even depressed [16, 17]. Some people may encounter emotional challenges for the first time, while others may have pre-existing mental health disorders that are made worse. Emotional sensitivity can also be increased by the strain of adjusting to new lifestyle demands as well as the environmental and social factors connected to severe weight reduction [18, 19]. It's imperative to recognize and manage mood swings and irritability in bariatric patients since they can impact the patient's quality of life, long-term weight control success, and compliance with postoperative instructions, in addition to their mental health.

Although bariatric surgery is an effective intervention for long-term weight loss, limited attention has been given to its psychological consequences, particularly mood swings and irritability, which may significantly affect postoperative adjustment and outcomes. Existing literature largely focuses on physical outcomes, while evidence on emotional and behavioral changes, especially in relation to hormonal shifts, nutritional deficiencies, and psychosocial adaptation, remains insufficient. This study aims to determine the prevalence and contributing factors of irritability and mood swings in post-bariatric surgery patients.

METHODS

This prospective observational study was carried out in the International Metabolic and Bariatric Center, Afridi Medical Complex, Peshawar, over a year, from February 2022 to January 2023. Before enrolment, all subjects were given their informed permission, and the Institutional Review Board (IRB) granted ethical approval with reference no.234/DME/AMC/2022. A sample size of 1014 patients was

determined by using the WHO sample size calculator. The calculation was based on a 95% confidence interval (CI), 5% margin of error (d), population (50%), and estimated prevalence (p) of 85.8%. The sample size was calculated using the formula [20]: $n = Z^2 \times P \times (1-P) / d^2$. A non-probability consecutive sampling technique was applied in this study. Patients who had bariatric surgery procedures such as Sleeve gastrectomy, Roux-en-Y gastric bypass (RYGB), and micro gastric bypass were among the surgeries carried out. Patients between the ages of 18-65 years, of both genders, who had no history of drug abuse or had any psychiatric conditions that required medication were eligible. Patients who refused to participate in follow-up or had any unstable medical conditions were excluded. Further, patients below 18 and above 65 years were not included in the current study. Profile of Mood States (POMS), a 5-point Likert scale comprised of 65 65-item inventory with six different subscales, including tension-anxiety, depression-dejection, Anger-Hostility, Fatigue-Inertia, Confusion-Bewilderment and Vigour-Activity. The scoring of this scale ranges from ('0=not at all' to '4=very strong') and is widely used for the assessment of mood [21]. The Irritation Questionnaire (IRQ), a two 4-point Likert scale, was employed to assess the frequency and severity of irritation symptoms. For frequency, the score ranges from ('0=never' to '3=most of the time') and for severity of symptoms, the score ranges from ('0=not at all' to '3=very much so') [22]. The Postoperative Adaptation Questionnaire (PAQ), a 15-item questionnaire, is used to assess how well the patient has adapted to lifestyle modifications, including eating habits and body image. Each item is reported on a 5-point Likert scale ranging from (0=strongly disagree to 5=strongly agree). Further sleep disorders, which can be a factor in mood swings, were assessed through the Sleep Quality Index (SQI). SQI is an 8-item questionnaire with three categories weighted 0, 1, and 2 and scoring varied from 0-16 (23). Blood samples were collected after overnight fasting, centrifuged, and serum was stored at -80°C, at baseline (pre-operative), 3rd month, and 6th month. Vitamin B12 and D levels were measured using chemiluminescent immunoassays, a highly sensitive, specific and suitable laboratory technique for detecting low serum concentrations of these micronutrients. Serotonin, ghrelin, and leptin were evaluated using ELISA kits (Elabscience, USA) following the manufacturer's protocols. Clinical and Demographic Information that including Preoperative psychological history, age, gender, BMI, operation type, concomitant conditions, and other baseline variables, were taken from medical records. Before the surgery, patients were evaluated to determine baseline levels of hormones such as serotonin, ghrelin and leptin, which are associated with appetite regulation. Nutritional health was assessed via blood levels of vitamin B12 and vitamin D. Mood and irritability were evaluated through valid tools such as Profile of Mood States (POMS)

and Irritability Questionnaire (IRQ). Participants filled out the IRQ and POMS questionnaires at follow-up assessments that were held three and six months after surgery. To track variations in hormone and nutritional levels, blood samples were taken at regular intervals. Samples were then centrifuged to separate serum and then stored at -80°C for preservation. The PAQ and SQI were used to collect psychosocial and sleep-related data during follow-ups. SPSS version 26.0 was used to analyze the data. Demographic information and baseline traits were compiled using descriptive statistics. Categorical variables such as gender, nutritional deficiencies, and surgery type were represented in percentages and frequencies, while the data in numerical form, such as age, BMI, irritability, mood disturbances, and nutritional and hormonal changes, were illustrated in mean and standard deviation. Relationships between mood swings and dietary or hormonal factors were investigated using an independent sample t-test, and the impact on hormonal and nutritional changes was found by multiple regression, and $p \leq 0.05$ was chosen as the threshold for statistical significance.

RESULTS

The study comprised 1,014 patients, whose mean age was 33 ± 13 years. Of those who took part, 659 (65%) were men and 335 (35%) were women. Before surgery, the average BMI was $29 \pm \text{kg/m}^2$. The majority of individuals had mini gastric bypass 385 (38%), followed by sleeve gastrectomy 304 (32%) and Roux-en-Y gastric bypass (RYGB) 324 (30%). According to the Profile of Mood States (POMS), 679 (67%) of participants had clinically evident mood abnormalities before surgery, and 740 (73%) of participants experienced mild irritation. With 284 (28%) of patients lacking vitamin B12 and 730 (72%) lacking vitamin D, nutritional deficiencies were common (Table 1).

Table 1: Baseline Demographic and Clinical Characteristics

Variables	Categories	Mean \pm SD	Frequency (%)
Mean Age (Years)	18-65	33 ± 13	—
Gender	Female	-	(335) 35%
	Male		659 (65%)
Mean BMI (kg/m^2)	-	29 ± 3	—
Surgery Type (%)	Sleeve Gastrectomy	-	385 (38%)
	Roux-en-Y Gastric		304 (30%)
	Mini Gastric Bypass		324 (32%)
Nutritional Deficiencies	Vitamin B12	-	284 (28%)
	Vitamin D		730 (72%)

Assessments conducted after surgery showed a marked rise in irritation. Compared to 740 (73%) before surgery, 902 (89%) of patients experienced moderate to severe irritation at three months ($p < 0.001$). After six months, this percentage dropped to 446 (44%). Increased frustration, trouble controlling emotions, and heightened

susceptibility to small stimuli were all signs of irritability. The percentage of patients reporting clinically significant mood disturbances increased from 679 (67%) before surgery to 844 (83%) at three months and 355 (35%) at six months ($p < 0.001$), indicating a significant worsening of mood swings following surgery a subsequent decrease after 6 months (Table 2).

Table 2: Changes in Mood and Irritability Scores Over Time

Measures	Baseline	3 rd Month	6 th Month	p-value
	n (%)			
Moderate to Severe Irritability	740 (73%)	902 (89%)	446 (44%)	<0.001
Mood Disturbances	679 (67%)	844 (83%)	355 (35%)	<0.001

After surgery, there were notable hormonal changes that were linked to mood swings and increased irritation. From 593 ± 31 ng/L ng/mL before surgery to 258 ± 18 ng/L at six months, ghrelin levels significantly decreased ($p < 0.001$), and patients who saw larger decreases also reported higher irritation scores ($p < 0.001$). At six months, serotonin levels dropped dramatically from 120 ± 20 ng/mL to 80 ± 15 ng/mL ($p = 0.04$), which led to more frequent mood swings. A subgroup of patients continued to have nutritional deficiencies; at six months, 345 (34%) were still lacking in vitamin B12 and 588 (58%) in vitamin D. Increased irritability and mood disorders were highly associated with these inadequacies (respectively, $p < 0.003$) (Table 3).

Table 3: Hormonal and Nutritional Changes

Measures	Baseline	6 Months	p-value	95% CI (Confidence Interval)
Serotonin (ng/mL)	120 ± 20	80 ± 15	<0.001	-12.7 to -8.3
Ghrelin (ng/L)	593 ± 31	258 ± 18	<0.001	-352.0 to -282.0
Vitamin B12 (ng/mL)	284 (28%)	345 (34%)	<0.003	0.12 to 0.25
Vitamin D (ng/mL)	730 (72%)	588 (58%)	<0.003	0.10 to 0.23

A decline in Postoperative Adaptation Questionnaire (PAQ) scores suggested that 345 (34%) of patients had postoperative adaptation issues. These patients showed noticeably more mood swings and irritation ($p < 0.001$). At six months, 304 (30%) of patients reported sleep difficulties, as judged by the Sleep Quality Index (SQI), compared to 65% before surgery ($p < 0.01$) (Table 4).

Table 4: Postoperative Adaptation Issues

Parameters	Preoperative	6 Months Post-operative	p-value	95% CI
Adaptation Issues	-	345 (34%)	<0.001	0.30 to 0.45
Mood Swings/Irritation	-	Increased	<0.001	0.50 to 0.60
Sleep Difficulties	659 (65%)	304 (30%)	<0.010	-0.25 to -0.15

DISCUSSION

The objectives of this study included the evaluation of the bariatric surgery candidates and post-surgical mood disorders, irritability, and postoperative adaptation over six months on a psychological, hormonal, and nutritional level.

Our results show alterations in mood and temper, which were at their worst at the three months following surgery and were relatively better at the six months. These changes are associated with a lower level of ghrelin and serotonin, and low levels of vitamin B12 and D remain low even many years postoperatively. Furthermore, only one-third of the patients responded that they did not have troubles with adaptation after surgery, and 304 (30%) reported persistent sleep disturbances. The results of this study, therefore, emphasize the importance of holistic care once the surgery is done. Our findings align with several previous studies, though with some differences in the magnitude and timing of changes. These changes of mood disturbances were described in bariatric patients by Loh et al., where 45% of the patients had depressive symptoms and 58% had anxiety symptoms before the surgery and changes which seem to increase in the first 3 months of the surgery, as it is seen in this study [24]. Ivezaj et al., also identified greater mood lability at three months after surgery, 13.1% of patients complained of irritability and depressive feelings. However, they noted that the resolution of these symptoms was slower, with 13 patients affected at 12 months in this study; only 44% of the patients still reported irritability at discharge at six months [25]. The changes in irritability and mood swings observed in bariatric surgery patients shortly after the surgery may be hormonal, nutritional, and psychosocial. One of the factors that aggravate such early mood disturbances is the hormonal change that follows surgery. Bariatric surgery results in a significant change in gut hormones, more specifically ghrelin, which is reported to fall swiftly after the operation. Ghrelin, which is also termed as the "hunger hormone", has some function in hunger as well as in the mood [26]. Hormonal fluctuation has been reported to affect brain reward and emotional centers, hence reducing or increasing irritability and poor emotional control [27]. In addition, serotonin, which plays a role in mood regulation, also decreases after surgery, making irritability and depressive-like symptoms worse [28]. Some of the changes also include the effects of nutritional deficiency, such as vitamin B12 and vitamin D deficiencies, which are usual after bariatric procedures, and worsen mood swings. These deficiencies may negatively affect the nervous system, which causes symptoms such as irritability, confusion, and fatigue [29]. Other aspects of the lifestyle change are expressed in psychosocial areas requiring interests, higher levels of stress, and changes in the parameters of the body, causing postoperative irritability [30]. Chronologically, these hormonal changes are balanced, nutritional deficiencies assuming a new diet, psychosocial adaptation is enhanced, and there is a decrease in mood disorders by six months after the surgery [31]. Therefore, the early postoperative period is a state of various physiological and psychological changes with relative amelioration of the process in the subsequent days

after surgery. The hormonal changes documented by Hamad et al., included a decrease in serotonin to an average of 54% of the preoperative levels, with a range of 36%–80% of the baseline value [32]. For that reason, Pasi et al., ascertained that 24.7% of persons who have undergone bariatric surgery experienced a decrease in serotonin levels, which are associated with increased irritability and mood swings, similar to our findings [33]. Our study observed that the percentage of patients who were deficient in vitamin B12 was 34% and those who were deficient in vitamin D was 58% at six months. These findings concur with those of Çalapkorur and Küçükkatirci, who reported that, among the bariatric surgery patients, 24.7% saw their levels of serotonin drop, which the authors attributed to increased irritability and mood swings, exactly as documented in this study [34]. Song and Fernstrom also observed that these vitamins were claimed to show some deficiency that is coupled with symptoms related to psychological side effects, such as irritation and depression [35]. In the present paper, we noted that 34% of patients experienced some problems with postoperative adjustment and reported more pronounced oscillations of mood and increased irritability. This is supported by Dalrymple et al., who observed that only 30% of bariatric patients had difficulties with emotional and psychological adjustment after the surgery. These difficulties were more pronounced in our poor sleep quality post-operative patients, who had 75% of them complaining of sleep disturbances at one-year follow-up, similar to our study, where 28% of the patients complained of sleep problems at six months follow-up [36]. Similarly, Lodewijks et al., also found that self-reported Sleep problems were still present in 30% of patients after six months of bariatric surgery and pointed out the probability of endocrine hormonal changes together with behavioural and lifestyle alterations [37]. Our assessment of sleep quality indicated that poor sleep quality changed from 65% before surgery to 30% at the six-month follow-up. By the above observation, Ab et al., reported that the patient had shown a 32% increase in sleep quality after surgery [38]. In a study by Sillo et al., they showed that 88.5% of bariatric patients had considerable clinical amendments in sleep apnea and other sleep disorders following weight loss, which is in agreement with our result of enhanced sleep quality; however, it also reveals that several patients continue to have some residual periods of sleep problem even after surgery [39]. This study is limited by its relatively short follow-up period of six months, which may not fully capture long-term psychological and nutritional outcomes after bariatric surgery. Additionally, being a single-center study may limit the generalizability of findings to broader bariatric populations. Future multicenter longitudinal studies with extended follow-up are recommended to better understand long-term psychological adaptation, hormonal

recovery, and nutritional deficiencies after bariatric surgery, and to develop structured post-operative mental health support programs.

CONCLUSIONS

The research found significant psychological and physiological changes after bariatric surgery. Mood disturbance and irritability status were increased at 3 months (83% and 89%, $p < 0.001$), and then improved by 6 months (35% and 44%, $p < 0.001$). Vitamins B12 and D deficiencies remained prevalent in 34% and 58% of patients throughout the 6-month follow-up, associated with increased irritability and mood issues ($p < 0.003$). Ghrelin and serotonin levels dropped significantly after surgery, contributing to mood changes ($p < 0.001$). Furthermore, Postoperative adaptation difficulties were observed in 34% of patients (PAQ), and sleep difficulties significantly reduced from 65% preoperatively to 30% at 6 months ($p < 0.010$).

Authors' Contribution

Conceptualization: MA¹

Methodology: MA¹, MA², WA

Formal analysis: MA², WA

Writing and Drafting: MA¹, MNDK, AHS

Review and Editing: MA¹, MA², WA, MNDK, AHS

All authors approved the final manuscript and take responsibility for the integrity of the work

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Aderinto N, Olatunji G, Kokori E, Olaniyi P, Isarinade T, Yusuf IA. Recent advances in bariatric surgery: a narrative review of weight loss procedures. *Annals of Medicine and Surgery*.2023Dec;85(12):6091-104.doi: 10.1097/MS9.0000000000001472.
- [2] Jumbe S, Bartlett C, Jumbe SL, Meyrick J. The effectiveness of bariatric surgery on long term psychosocial quality of life-A systematic review. *Obesity Research and Clinical Practice*.2016May; 10(3): 225-42. doi: 10.1016/j.orcp.2015.11.009.
- [3] Berger H, Ordemann J, Elbelt U, Hofmann T, Menenakos C, Berger H et al. Postoperative Management of Bariatric Surgery Patients. In *Obesity and Metabolic Surgery*. Berlin, Heidelberg: Springer Berlin Heidelberg.2022Oct:139-164.doi:10.1007/978-3-662-63227-7_14.
- [4] Brown RM, Guerrero-Hreins E, Brown WA, le Roux CW, Sumithran P. Potential gut-brain mechanisms behind adverse mental health outcomes of bariatric surgery. *Nature Reviews Endocrinology*.2021Sep;17(9):549-59. doi: 10.1038/s41574-021-00520-2.
- [5] Papamargaritis D and le Roux CW. Do gut hormones contribute to weight loss and glycaemic outcomes after bariatric surgery? *Nutrients*.2021Feb; 13(3): 762 .doi: 10.3390/nu13030762.
- [6] Barrea L, Salzano C, Pugliese G, Laudisio D, Frias-Toral E, Savastano S et al. The challenge of weight loss maintenance in obesity: a review of the evidence on the best strategies available. *International Journal of Food Sciences and Nutrition*.2022Nov; 73(8): 1030-46. doi: 10.1080/09637486.2022.2130186.
- [7] Makaronidis JM and Batterham RL. The role of gut hormones in the pathogenesis and management of obesity. *Current Opinion in Physiology*.2019 Dec;12: 1-1. doi: 10.1016/j.cophys.2019.04.007.
- [8] Moffett RC, Docherty NG, le Roux CW. The altered enteroendocrine repertoire following roux-en-Y-gastric bypass as an effector of weight loss and improved glycaemic control. *Appetite*.2021Jan;156: 104807. doi: 10.1016/j.appet.2020.104807.
- [9] Bhat N, Bhat J, Bhat K, Junaid I, Bhat A, Bhat J. Restoring gut biome balance for weight loss: clinical applications. In *Microbiome, Immunity, Digestive Health and Nutrition*.2022Jan:167-184.doi:10.1016/B978-0-12-822238-6.00022-4.
- [10] Stone LA, Harmatz ES, Goosens KA. Ghrelin as a stress hormone: implications for psychiatric illness. *Biological Psychiatry*.2020Oct;88(7):531-40.doi:10.1016/j.biopsych.2020.05.013.
- [11] Friedman JM. Leptin and the endocrine control of energy balance. *Nature Metabolism*.2019 Aug; 1(8): 754-64. doi: 10.1038/s42255-019-0095-y.
- [12] Platzer M, Fellendorf FT, Bengesser SA, Birner A, Dalkner N, Hamm C et al. The relationship between food craving, appetite-related hormones and clinical parameters in bipolar disorder. *Nutrients*.2020Dec; 13(1):76. doi:10.3390/nu13010076.
- [13] Hoepner CT, McIntyre RS, Papakostas GI. Impact of supplementation and nutritional interventions on pathogenic processes of mood disorders: A Review of the Evidence. *Nutrients*.2021Feb; 13(3): 767. doi: 10.3390/nu13030767.
- [14] Berthou C, Iliou JP, Barba D. Iron, neuro-bioavailability and depression. *EJHaem*.2022Feb; 3(1): 263-75. doi: 10.1002/jha2.321.
- [15] Noah L, Dye L, Bois De Fer B, Mazur A, Pickering G, Pouteau E. Effect of magnesium and vitamin B6 supplementation on mental health and quality of life in stressed healthy adults: Post-hoc analysis of a randomised controlled trial. *Stress and Health*.2021 Dec; 37(5): 1000-9. doi: 10.1002/smi.3051.

- [16] Renteria K, Nguyen H, Koh GY. The role of vitamin D in depression and anxiety disorders: A review of the literature. *Nutritional Neuroscience*. 2024 Mar; 27(3): 262-70. doi: 10.1080/1028415X.2023.2186318.
- [17] Conceição EM, Fernandes M, de Lourdes M, Pinto-Bastos A, Vaz AR, Ramalho S. Perceived social support before and after bariatric surgery: association with depression, problematic eating behaviors, and weight outcomes. *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity*. 2020 Jun; 25: 679-92. doi: 10.1007/s40519-019-00671-2.
- [18] Francois Z and Rizvi A. Psychiatric Complications of Bariatric Surgery. In *Stat Pearls* [Internet]. 2024 May. Stat Pearls Publishing.
- [19] Berg A. Untold stories of living with a bariatric body: long-term experiences of weight-loss surgery. *Sociology of Health and Illness*. 2020 Feb; 42(2): 217-31. doi: 10.1111/1467-9566.12999.
- [20] Naing L, Nordin RB, Abdul Rahman H, Naing YT. Sample size calculation for prevalence studies using Scalex and ScalaR calculators. *BioMed Central Medical Research Methodology*. 2022 Jul; 22(1): 209. doi: 10.1186/s12874-022-01694-7.
- [21] Petrowski K, Albani C, Zenger M, Brähler E, Schmalbach B. Revised short screening version of the profile of mood states (POMS) from the German general population. *Frontiers in Psychology*. 2021 May; 12: 631668. doi: 10.3389/fpsyg.2021.631668.
- [22] Craig KJ, Hietanen H, Markova IS, Berrios GE. The Irritability Questionnaire: a new scale for the measurement of irritability. *Psychiatry Research*. 2008 Jun; 159(3): 367-75. doi: 10.1016/j.psychres.2007.03.002.
- [23] Urponen H, Partinen M, Vuori I, Hasan J. Sleep quality and health: Description of the sleep quality index. In *Sleep and Health Risk*. Berlin, Heidelberg: Springer Berlin Heidelberg. 1991: 555-558. doi: 10.1007/978-3-642-76034-1_63.
- [24] Loh HH, Francis B, Lim LL, Lim QH, Yee A, Loh HS. Improvement in mood symptoms after post-bariatric surgery among people with obesity: A systematic review and meta-analysis. *Diabetes/Metabolism Research and Reviews*. 2021 Nov; 37(8): e3458. doi: 10.1002/dmrr.3458.
- [25] Ivezaj V and Grilo CM. When mood worsens after gastric bypass surgery: characterization of bariatric patients with increases in depressive symptoms following surgery. *Obesity Surgery*. 2015 Mar; 25(3): 423-9. doi: 10.1007/s11695-014-1402-z.
- [26] Geloneze B, Tambascia MA, Pilla VF, Geloneze SR, Repetto EM, Pareja JC. Ghrelin: a gut-brain hormone: effect of gastric bypass surgery. *Obesity Surgery*. 2003 Feb; 13(1): 17-22. doi: 10.1381/096089203321136539.
- [27] Maric I, López-Ferreras L, Bhat Y, Asker M, Borchers S, Bellfy L et al. From the stomach to locus coeruleus: new neural substrate for ghrelin's effects on ingestive, motivated and anxiety-like behaviors. *Frontiers in Pharmacology*. 2023 Nov; 14: 1286805. doi: 10.3389/fphar.2023.1286805.
- [28] Bouchard-Mercier A, de Toro-Martín J, Nadeau M, Lescelleur O, Lebel S, Richard D et al. Molecular remodeling of adipose tissue is associated with metabolic recovery after weight loss surgery. *Journal of Translational Medicine*. 2022 Jun; 20(1): 283. doi: 10.1186/s12967-022-03485-6.
- [29] Argyrakopoulou G, Konstantinidou SK, Dalamaga M, Kokkinos A. Nutritional deficiencies before and after bariatric surgery: prevention and treatment. *Current Nutrition Reports*. 2022 Jun; 11(2): 95-101. doi: 10.1007/s13668-022-00400-9.
- [30] Sarwer DB, Wadden TA, Fabricatore AN. Psychosocial and behavioral aspects of bariatric surgery. *Obesity Research*. 2005 Apr; 13(4): 639-48. doi: 10.1038/oby.2005.71.
- [31] O'Brien PE, Hindle A, Brennan L, Skinner S, Burton P, Smith A et al. Long-term outcomes after bariatric surgery: a systematic review and meta-analysis of weight loss at 10 or more years for all bariatric procedures and a single-centre review of 20-year outcomes after adjustable gastric banding. *Obesity Surgery*. 2019 Jan; 29: 3-14. doi: 10.1007/s11695-018-3525-0.
- [32] Hamad GG, Helsel JC, Perel JM, Kozak GM, McShea MC, Hughes C et al. The effect of gastric bypass on the pharmacokinetics of serotonin reuptake inhibitors. *American Journal of Psychiatry*. 2012 Mar; 169(3): 256-63. doi: 10.1176/appi.ajp.2011.11050719.
- [33] Pasi P, Kröll D, Siegfried A, Sykora M, Wildisen A, Milone C et al. Plasma concentrations of SSRI/SNRI after bariatric surgery and the effects on depressive symptoms. *Frontiers in Psychiatry*. 2023 Apr; 14: 1132112. doi: 10.3389/fpsyg.2023.1132112.
- [34] Çalapkörür S and Küçükkatirci H. Vitamin deficiencies and prevention methods after bariatric surgery. *Mini-Invasive Surgery*. 2020 Mar; 4: N-A. doi: 10.20517/2574-1225.2019.51.
- [35] Song A and Fernstrom MH. Nutritional and psychological considerations after bariatric surgery. *Aesthetic Surgery Journal*. 2008 Mar; 28(2): 195-9. doi: 10.1016/j.asj.2008.01.005.
- [36] Dalrymple KL, Maleva V, Chelminski I, Zimmerman M. Judgment towards emotions as a mediator of the relationship between emotional eating and depression symptoms in bariatric surgery candidates. *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity*. 2022 Dec; 27(8): 3675-83. doi: 10

- .1007/s40519-022-01508-1.
- [37] Lodewijks Y, Schonck F, Nienhuijs S. Sleep quality before and after bariatric surgery. *Obesity Surgery*. 2023 Jan; 33(1):279-83. doi:10.1007/s11695-022-06387-0.
- [38] Ab Majid NL, Vanoh D, Zainuddin NZ, Hashim MN. Post bariatric surgery complications, nutritional and psychological status. *Asia Pacific Journal of Clinical Nutrition*. 2024 Jun; 33(2): 162.
- [39] Sillo TO, Lloyd-Owen S, White E, Abolghasemi-Malekabadi K, Lock-Pullan P, Ali M *et al.* The impact of bariatric surgery on the resolution of obstructive sleep apnoea. *BioMed Central Research Notes*. 2018 Dec; 11: 1-6. doi: 10.1186/s13104-018-3484-5.