



Original Article

Correlation between Vitamin D Deficiency and Osteopenia in Women Residing in Karachi, Pakistan

Yameen Bocha¹, Anita Haroon², Benish Zafar³, Kahkashan Perveen³, Naheed Akhtar⁴, Padma Rathore⁵ and Saher Mushtaque⁶¹Department of Medicine, Sindh Govt Lyari General Hospital, Karachi, Pakistan²Department of Medicine, Hamdard University Hospital, Karachi, Pakistan³Department: Biochemistry, Baqai Medical University, Karachi, Pakistan⁴Department of Anatomy, Karachi Institute of Medical Sciences, Combined Military Hospital, Karachi, Pakistan⁵Department of Physiology, Jinnah Sindh Medical University, Karachi, Pakistan⁶Department of Community Dentistry, Sir Syed College of Medical Sciences, Karachi, Pakistan

ARTICLE INFO

Keywords:

Osteopenia, Pakistan, Women, Vitamin D

How to Cite:

Bocha, Y., Haroon, A., Zafar, B., Perveen, K., Akhtar, N., Rathore, P., & Mushtaque, S. (2025). Correlation between Vitamin D Deficiency and Osteopenia in Women Residing in Karachi, Pakistan : Vitamin D Deficiency and Osteopenia in Women . Pakistan Journal of Health Sciences, 6(6), 232-236. <https://doi.org/10.54393/pjhs.v6i6.3073>

*Corresponding Author:

Saher Mushtaque

Department of Community Dentistry, Sir Syed College of Medical Sciences, Karachi, Pakistan
dr.saher83@gmail.com

Received Date: 15th April, 2025Revised Date: 16th June, 2025Acceptance Date: 20th June, 2025Published Date: 30th June, 2025

ABSTRACT

Osteopenia is one of the primary health concerns among Pakistani women. **Objective:** To assess the Correlation Between Vitamin D Deficiency and Osteopenia in Women in Karachi, Pakistan. **Methods:** A cross-sectional survey was conducted among 200 women in Karachi, Pakistan. A self-reported questionnaire was given to all participants so that they could fill in the demographic data. Only participants who were willing to participate in the study were included, and those who failed to provide any data regarding the status of menopause were excluded. SPSS version 23.0 was used to analyze the data. **Results:** The 200 women in the study had a high reproductive burden (mean pregnancies = 12), with a mean age of 65.2 years and a BMI of 25.6 kg/m². Age-related decreases in bone mineral density were significant ($p=0.0002$), and osteoporosis was most prevalent in older age groups. The distribution of vitamin D deficiency across age groups was not statistically significant ($p=0.22$), but it was associated with ageing ($p=0.011$). **Conclusions:** It was concluded that although vitamin D deficiency is common among women, age is a better indicator of deteriorating bone mineral density. Age-focused bone health interventions are needed, as osteoporosis and osteopenia were significantly correlated with advancing age.

INTRODUCTION

The long-term health of the ageing population is being negatively impacted by osteoporosis and osteopenia, which raises the risk of death, reduces autonomy, increases hospitalization rates, and places a significant financial and social load on the healthcare system. The longer life expectancy is linked to a higher prevalence of age-related illnesses, including osteopenia [1]. Osteopenia is one of the biggest health issues faced by Pakistani women currently [2]. The World Health Organization (WHO) defines osteopenia as having a T-score between -1.0 and -

2.5, which indicates that bone density is lower than usual but not low enough to be categorized as osteoporosis [3]. A T-score of -2.5 or less indicates osteoporosis, which is characterized by a more severe loss of bone mass [4]. A degenerative disease that affects all of the bones in the body is called osteoporosis [5]. When the levels of bone mass density are below normal, it is called osteopenia [6]. Osteopenia is 40.40% and osteoporosis is 19.75% prevalent worldwide [7]. Hence, the burden of these conditions has been rising for almost three decades [8]. It

has a complex etiology since numerous factors affect bone health, such as exercise and diet [9, 10]. Pakistani women's osteopenia rates are high due to sedentary lifestyles, a deficiency in foods high in calcium, and the popularity of fast food. Osteopenia is primarily linked to inadequate vitamin D intake or lack of availability of sufficient sunlight [11]. Serum levels of 25-hydroxyvitamin D (25 (OH)D) are commonly used to determine vitamin D status, and classifications have become more uniform in recent years. Gupta et al., 2022 define vitamin D deficiency as serum 25 (OH)D levels below 12 ng/mL (30 nmol/L), insufficiency as levels between 12 and 20 ng/mL (30 and 50 nmol/L), and sufficiency as levels above 20 ng/mL (50 nmol/L). These cutoff points are used for screening and treatment decisions in both adult and pediatric populations, and they are based on correlations with outcomes related to bone health [12]. The main causes of vitamin D deficiency are nutritional deficiencies, limited sun exposure among women, and other chronic conditions. Appropriate amounts of vitamin D are available in the subcontinent with adequate sun exposure. The body uses vitamin D as a hormone, and it is crucial for the mineralization of bones. It also supports immunological function, muscular integrity, and cell proliferation. It contributes to healthy skin development, reproduction, and other metabolic processes. Given its increasing significance, scientists advise taking it at a dosage of at least 1000 IU. Worldwide, the prevalence of vitamin D deficiency is estimated to be 50%. Obesity and malnutrition can be detrimental to one's health and can cause bone problems such as osteoporosis, osteoarthritis, sarcoidosis-related osteoporosis, and osteopenia. Food, medication, and nutritional supplements are used to treat metabolic bone disease [13]. Pakistan is underfunded to address this disease's burden because of its slow progression and high cost of care. A population-based study found that osteopenia and osteoporosis affect millions of people in Pakistan [14]. The Osteoporosis Risk Assessment Instrument now includes current estrogen therapy in addition to the Osteoporosis Self-Assessment Tool and OST for people residing on the Asia continent. Hence, the prognostic instruments (OST, OSTA, and Osteoporosis Index) of Risk, as well as straightforward computed osteoporosis risk estimation. Concentrate on postmenopausal women's osteoporosis risk. Certain biomarkers, such as blood or urine indicators, are readily available during normal medical examinations and have been shown to enhance the sensitivity of osteoporosis prediction at the preclinical stage [15]. This study aims to assess the correlation between vitamin D deficiency and osteopenia in women residing in Karachi, Pakistan

METHODS

In Karachi, Pakistan, 200 participants with an average age of 65.2 years were recruited for this cross-sectional study between June and September 2024. With $Z=1.96$ at a 95% confidence level, p =expected prevalence, and d =margin of error (0.05), the sample size was calculated using open epi based on past research conducted by Khan et al., [16] showing high rates among Karachi women (an expected prevalence of 75% for vitamin D deficiency. We recruited 200 participants to account for the estimated sample size of 196 with an 80% power and a 5% margin of error. A non-probability purposive sampling strategy was used to find participants from outreach clinics and community health centers located throughout Karachi. Women who were at least 45 years old, Karachi residents, and willing to give informed consent were among the requirements for inclusion. Women who reported not being menopausal or who were receiving treatment for any bone-related conditions or recent fractures were not included in the study. To further reduce confounding, people taking vitamin D supplements or having comorbid conditions like liver or kidney disease were not included. The Sir Syed Medical and Dental College for Girls' Ethical Review Committee granted ethical approval for the study (Ref. No: 032 SSCMS-Ethics/2024). After being made aware of the goals and methods of the study, each participant gave their written informed consent. A structured, self-reported questionnaire that was pilot-tested and approved by public health specialists for relevance and clarity was used to gather data. Age, weight, height, number of pregnancies, lifestyle factors, history of vitamin D supplementation, sun exposure, and known comorbidities were all included in the questionnaire. Where possible, laboratory studies were used to supplement clinical data. Dual-Energy X-ray Absorptiometry (DEXA) scans were used to measure bone mineral density (BMD). According to WHO guidelines, osteoporosis was defined as a T-score ≤ -2.5 , and osteopenia as a T-score between -1.0 and -2.5 . According to Gupta et al., vitamin D status is classified as deficient (<12 ng/mL), insufficient (12–20 ng/mL), and sufficient (>20 ng/mL) based on serum 25-hydroxyvitamin D [(5(OH)D) levels [12]. SPSS version 23.0 was used for statistical analysis. Standard deviations and means were used to summarize continuous variables. For non-parametric data, independent samples were compared using the Mann-Whitney U test. Since vitamin D status and BMD were both ordinal/non-parametric variables, the relationship between them was evaluated using Spearman's rank correlation coefficient. Unless otherwise noted, all tests were two-tailed and performed at a 5% level of significance. Correlation analyses took into account confounding variables like age and BMI.

RESULTS

The results of the study revealed that the demographic characteristics of the participants, showing an average age of 65.2 years, mean BMI of 25.6 kg/m² (overweight range), and a high average number of pregnancies (12), reflecting a population at elevated risk for bone health issues (Table 1).

Table 1: Participants' Demographic Details

Variables	Mean
Age (Years)	65.2
Weight (kg)	59
Height (m)	1.51
BMI (kg/m ²)	25.6
No. of Pregnancies	12

The mean age of participants is categorized by vitamin D classification and bone mineral density (BMD) status. The mean age of the 200 participants was 67.3 ± 9.3 years for those with normal BMD (n=28), whereas the mean ages of those with osteopenia (n=117) and osteoporosis (n=55) were higher at 68.5 ± 8.6 years and 74.2 ± 9.9 years, respectively. Growing older is linked to a higher risk of having less bone mineral density, as evidenced by the statistically significant difference in mean age between the BMD categories (p=0.0002). Similarly, participants with vitamin D deficiency (n=74) had the highest mean age of 71.5 ± 9.4 years, followed by those with sufficient levels (n=16; 69.3 ± 9.1 years) and those with insufficient levels (n=110; 68.6 ± 8.5 years). Additionally, this difference was statistically significant (p=0.011), pointing to a possible connection between deteriorating vitamin D levels and ageing. These results highlight the link between vitamin D deficiency, ageing, and deteriorating bone health in the population under study (Table 2).

Table 2: The Average Patient Age (in Years) for Osteopenia and Vitamin D3

Variables	n	Age Mean ± SD	p-Value
Classification of Bone Mineral Density			
Normal	28	67.3 ± 9.3	0.0002
Osteopenia	117	68.5 ± 8.6	
Osteoporosis	55	74.2 ± 9.9	
Classification of Vitamin D			
Deficiency	74	71.5 ± 9.4	0.011
Insufficient	110	68.6 ± 8.5	
Sufficient	16	69.3 ± 9.1	

The mean age of participants is categorized by vitamin D classification and bone mineral density (BMD) status. The mean age of the 200 participants was 67.3 ± 9.3 years for those with normal BMD (n=28), whereas the mean ages of those with osteopenia (n=117) and osteoporosis (n=55) were higher at 68.5 ± 8.6 years and 74.2 ± 9.9 years, respectively.

Growing older is linked to a higher risk of having less bone mineral density, as evidenced by the statistically significant difference in mean age between the BMD categories (p=0.0002). Similarly, participants with vitamin D deficiency (n=74) had the highest mean age of 71.5 ± 9.4 years, followed by those with sufficient levels (n=16; 69.3 ± 9.1 years) and those with insufficient levels (n=110; 68.6 ± 8.5 years). Additionally, this difference was statistically significant (p=0.011), pointing to a possible connection between deteriorating vitamin D levels and ageing. These results highlight the link between vitamin D deficiency, ageing, and deteriorating bone health in the population under study (Table 2).

Table 3: Correlation of Women's Age with Each Variable

Variables	45-59 Years	60-69 Years	70-79 Years	80-99 Years	p- Value
Bone Mineral Density					
Normal	6(3%)	15(7%)	15(7%)	12(6%)	0.005
Osteopenia	9(4%)	30(15%)	31(15%)	19(9%)	
Osteoporosis	3(1%)	13(6%)	19(9%)	28(14%)	
Vitamin D Status					
Deficiency	10(5%)	21(10%)	27(13%)	15(7%)	0.22
Insufficient	16(8%)	30(15%)	30(15%)	14(7%)	
Sufficient	6(3%)	11(5%)	9(4%)	11(5%)	

DISCUSSION

Osteopenia is a symptom of osteoporosis and a major indicator of bone health among women in Asia [18]. Deficits in vitamin D, which is essential for calcium absorption, are uncommon in young, healthy people but common in elderly persons, particularly in areas with little sunshine [19]. The results of our investigation showed that while vitamin D deficiency is widespread, its association with osteopenia is complicated and is probably impacted by confounding variables such as age, body mass index, and reproductive history. Interestingly, the results indicated that there was no correlation between vitamin D levels and BMD status across age groups (p=0.22), even though 74 out of 200 people had vitamin D deficiency. The lack of a statistically significant correlation between vitamin D levels and BMD status (p=0.11) implies that while vitamin D may affect bone health, age may be a more relevant predictor of BMD. Women with osteoporosis had a considerably higher mean age (74.2 ± 9.9 years) than those with osteopenia (68.5 ± 8.6 years) and normal bone mineral density (67.3 ± 9.3 years), which supports the notion that ageing-related physiological changes have a major impact on bone density (p=0.0002). Deteriorating bone health may also result from the cumulative demands on calcium stores during pregnancy and lactation [20]. This problem is especially important in South Asia, where there are several lifestyle variables, limited sun exposure, and low consumption of foods high in vitamin D [21]. According to this study, women

with osteopenia and osteoporosis frequently have vitamin D deficiency, and age seems to have a bigger influence on their BMD status in this group. Future studies should consider longitudinal methods, looking at additional important elements including exercise, dietary patterns, and socio-economic status.

CONCLUSIONS

It was concluded that bone mineral density (BMD) significantly decreases with age, and osteopenia and vitamin D deficiency are highly prevalent among older women in Karachi. Despite being widespread, vitamin D deficiency did not statistically significantly correlate with BMD status, indicating that ageing is a more reliable indicator of declining bone health. To prevent bone-related complications in postmenopausal women, these findings emphasize the necessity of early screening and age-targeted interventions.

Authors Contribution

Conceptualization: YB, SM

Methodology: YB, KP, NA, SM

Formal analysis: BZ, SM

Writing review and editing: AH, KP, NA, PR

All authors have read and agreed to the published version of the manuscript

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Fan Y, Li Q, Liu Y, Miao J, Zhao T, Cai J et al. Sex-and Age-Specific Prevalence of Osteopenia and Osteoporosis: Sampling Survey. *Journal of Medical Internet Research Public Health and Surveillance*. 2024 Apr; 10(1): e48947. doi: 10.2196/48947.
- [2] Nazir S, Sagheer U, Riaz S, Khan ZM. Frequency of Osteoporosis and Osteopenia and its associated factors among general population in Faisalabad. *The Professional Medical Journal*. 2019 Dec; 26(12): 2216-21. doi: 10.29309/TPMJ/2019.26.12.4073.
- [3] Varacallo M, Seaman TJ, Jandu JS, Pizzutillo P. Osteopenia. 2023 Aug.
- [4] Vierucci F, Del Pistoia M, Fanos M, Erba P, Saggese G. Prevalence of hypovitaminosis D and predictors of vitamin D status in Italian healthy adolescents. *Italian Journal of Pediatrics*. 2014 Dec; 40: 1-9. doi: 10.1186/1824-7288-40-54.
- [5] Zhang L, Luo X, Liu H, Zhu W, Zhang X, Zhu S et al. Prevalence and risk factors of osteoporosis and osteopenia among residents in Hubei province, China. *Archives of Osteoporosis*. 2023 Apr; 18(1): 49. doi: 10.1007/s11657-023-01245-7.
- [6] Lin X, Guo H, Lian Y, Kou J, Wang G, Chen Y et al. Osteoporosis and related health status among the elderly urban residents in elderly-care inns in Beijing, a multicenter DXA survey. *Frontiers in Endocrinology*. 2022 Jul; 13: 875678. doi: 10.3389/fendo.2022.875678.
- [7] Xiao PL, Cui AY, Hsu CJ, Peng R, Jiang N, Xu XH et al. Global, regional prevalence, and risk factors of osteoporosis according to the World Health Organization diagnostic criteria: a systematic review and meta-analysis. *Osteoporosis International*. 2022 Oct; 33(10): 2137-53. doi: 10.1007/s00198-022-06454-3.
- [8] Shen Y, Huang X, Wu J, Lin X, Zhou X, Zhu Z et al. The global burden of osteoporosis, low bone mass, and its related fracture in 204 countries and territories, 1990-2019. *Frontiers in Endocrinology*. 2022 May; 13: 882241. doi: 10.3389/fendo.2022.882241.
- [9] De Sire A, Gallelli L, Marotta N, Lippi L, Fusco N, Calafiore D et al. Vitamin D deficiency in women with breast cancer: a correlation with osteoporosis? A machine learning approach with multiple factor analysis. *Nutrients*. 2022 Apr; 14(8): 1586. doi: 10.3390/nu14081586.
- [10] Grili PP, Vidigal CV, Cruz GF, Albergaria BH, Marques-Rocha JL, Pereira TS et al. Nutrient patterns and risk of osteopenia in postmenopausal women. *Nutrients*. 2023 Mar; 15(7): 1670. doi: 10.3390/nu15071670.
- [11] Chadgal S and Talwar I. Prevalence of osteopenia and osteoporosis among young adult girls residing in the hostels of Panjab University, Chandigarh. *Prevalence*. 2017 Nov; 2(6).
- [12] Gupta P, Dabas A, Seth A, Bhatia VL, Khadgawat R, Kumar P et al. Indian Academy of Pediatrics Revised (2021) guidelines on prevention and treatment of vitamin D deficiency and rickets. *Indian Pediatrics*. 2022 Feb; 59(2): 142-58. doi: 10.1007/s13312-022-2448-y.
- [13] Zahid S, Hanif A, Faridi TA, Masood Z, John A. Frequency of Osteopenia and its association with dietary factors among female population aged 18-60 years: Osteopenia and its association with dietary factors. *Diet Factor (Journal of Nutritional and Food Sciences)*. 2020; 01(02): 48-52. doi: 10.54393/df.v1i02.2.
- [14] De Martinis M, Allegra A, Sirufo MM, Tonacci A, Pioggia G, Raggiunti M et al. Vitamin D deficiency, osteoporosis and effect on autoimmune diseases and hematopoiesis: a review. *International Journal of*

- Molecular Sciences. 2021 Aug; 22(16): 8855. doi: 10.3390/ijms22168855.
- [15] Chen JH, Chen YC, Tsai MK, Chiou JM, Lee WC, Tsao CK, Tsai KS, Chie WC. Predicting the risk of osteopenia for women aged 40–55 years. *Journal of the Formosan Medical Association*. 2017 Nov; 116(11): 888–96. doi: 10.1016/j.jfma.2017.01.003.
- [16] Khan AH, Iqbal R, Naureen G, Dar FJ, Ahmed FN. Prevalence of vitamin D deficiency and its correlates: results of a community-based study conducted in Karachi, Pakistan. *Archives of Osteoporosis*. 2012 Dec; 7: 275–82. doi: 10.1007/s11657-012-0108-x.
- [17] Haq AA, Memon I, Mahmood N, Shah FR, Malik F. Prevalence of osteopenia and osteoporosis in women using dual energy x-ray absorptiometry scan. *Pakistan Journal of Physiology*. 2021 Jun; 17(2): 27–30. doi: 10.69656/pjp.v17i2.1316.
- [18] Tang G, Feng L, Pei Y, Gu Z, Chen T, Feng Z. Low BMI, blood calcium and vitamin D, kyphosis time, and outdoor activity time are independent risk factors for osteoporosis in postmenopausal women. *Frontiers in Endocrinology*. 2023 Oct; 14: 1154927. doi: 10.3389/fendo.2023.1154927.
- [19] van Schoor N, de Jongh R, Lips P. Worldwide vitamin D status. *Feldman and Pike's Vitamin D*. 2024 Jan: 47–75. doi: 10.1016/B978-0-323-91338-6.00004-5.
- [20] Min C, Yoo DM, Wee JH, Lee HJ, Choi HG. High-intensity physical activity with high serum vitamin D levels is associated with a low prevalence of osteopenia and osteoporosis: A population-based study. *Osteoporosis International*. 2021 May; 32: 883–91. doi: 10.1007/s00198-020-05746-w.
- [21] Shamsi U, Azam I, Shamsi A, Shamsi D, Callen D. Frequency and determinants of vitamin D deficiency among premenopausal and postmenopausal women in Karachi Pakistan. *BioMed Central Women's Health*. 2021 May; 21(1): 194. doi: 10.1186/s12905-021-01339-9.