



## Case Series

## Histopathological and Radiological Findings in Nasolabial Cysts

Vijay Kumar Khatri<sup>1</sup>, Syed Muhammad Moosa Raza<sup>2</sup>, Sabhagi Kumari<sup>2</sup>, Syed Mohsin Ahmed Rizvi<sup>2</sup>, Saima Batool Jaffri<sup>3</sup>, Fiza Irfan<sup>3</sup> and Ali Muntazir Naqvi<sup>4\*</sup><sup>1</sup>Department of ENT, Al-Tibri Medical College and Hospital, Karachi, Pakistan<sup>2</sup>Department of Oral Pathology, Sindh Institute of Oral Health Sciences, Jinnah Sindh Medical University, Karachi, Pakistan<sup>3</sup>Department of Oral Medicine and Diagnosis, Sindh Institute of Oral Health Sciences, Jinnah Sindh Medical University, Karachi, Pakistan<sup>4</sup>Prof. Dr. Muhammad Saleh Memon Research and Diagnostic Lab, Al Tibri Medical College and Hospital, Karachi, Pakistan

## ARTICLE INFO

**Keywords:**

Nasolabial Cysts, Histopathological Features, Radiological Features, Bone Remodelling

**How to Cite:**Khatri, V. K., Raza, S. M. M., Kumari, S., Rizvi, S. M. A., Jaffri, S. B., Irfan, F., & Naqvi, A. M. (2025). Histopathological and Radiological Findings in Nasolabial Cysts: Radiological Findings in Nasolabial Cysts. *Pakistan Journal of Health Sciences*, 6(6), 305-309. <https://doi.org/10.54393/pjhs.v6i6.3117>**\*Corresponding Author:**Ali Muntazir Naqvi  
Department of ENT, Al-Tibri Medical College and Hospital, Pakistan  
[ali.muntazir@gmail.com](mailto:ali.muntazir@gmail.com)Received Date: 30<sup>th</sup> April, 2025Revised Date: 15<sup>th</sup> June, 2025Acceptance Date: 20<sup>th</sup> June, 2025Published Date: 30<sup>th</sup> June, 2025

## ABSTRACT

Nasolabial cysts are relatively rare, non-odontogenic soft tissue cysts that are usually found in the nasolabial fold, just beneath the nasal ala. **Objectives:** To analyse the clinical, histopathological, and radiological features of nasolabial cysts and evaluate their association with symptom duration, cyst size, and bone remodelling. **Methods:** This retrospective case series study included 10 patients diagnosed with nasolabial cysts. Patient data, including age, gender, ethnicity, cyst size, location, and duration of symptoms, were collected. Radiological assessments focused on bone remodelling and scalloping through computed tomography (CT) scans. Histopathological examination of excised cysts assessed the type of epithelial lining. Statistical analysis was performed to evaluate associations between cyst characteristics, symptom duration, and radiological findings, using Pearson's correlation and Chi-square tests. **Results:** The mean age of patients was 44.9 years, with a male predominance (60%). A significant association was found between longer symptom duration and the presence of bone changes on CT ( $p=0.03$ ), and a moderate positive correlation existed between cyst size and symptom duration ( $r=0.32$ ,  $p=0.009$ ). **Conclusions:** It was concluded that nasolabial cysts can lead to bone remodelling when symptoms persist for extended periods. Early diagnosis and treatment are critical in preventing complications. Further research is needed to explore minimally invasive treatment approaches for managing nasolabial cysts.

## INTRODUCTION

Nasolabial cysts are relatively rare, non-odontogenic soft tissue cysts that are usually found in the nasolabial fold, just beneath the nasal ala. These cysts are also referred to as nasoalveolar or Klestadt's cysts in some literature. They are different from other common facial cysts because of their origin, which is believed to be from the remnants of the embryonic nasolacrimal duct [1]. Most cases are discovered incidentally, as many patients do not present with obvious symptoms. However, in some instances, individuals may develop visible swelling in the nasolabial region, experience nasal obstruction, or suffer from localized pain and infection. When symptoms do occur, they can be quite uncomfortable and may interfere with

daily activities or lead to cosmetic concerns that affect the patient's overall well-being [2]. In terms of demographic trends, these cysts are seen more often in women, particularly those in their middle age, which suggests a possible hormonal or developmental predisposition. Moreover, studies and clinical observations have shown that nasolabial cysts are more frequently encountered in Asian populations compared to others [1, 3]. Even though these cysts are benign and don't have malignant potential, their anatomical location close to vital facial structures can lead to both aesthetic issues and functional impairments if the cyst enlarges or becomes secondarily infected. From a histopathological point, the nasolabial cysts are usually



lined with pseudostratified columnar epithelium that contains mucus cells called goblet cells. In many cases, it can also be squamous metaplasia, but it does not happen very much [4]. This type of epithelial lining gives support to the theory that it is coming from the nasolacrimal duct. Accurate histology evaluation is very important to distinguish nasolabial cyst from other cyst lesions which also in the same area, like odontogenic or dermoid cyst and other non-odontogenic growings. This difference needs to be avoided to prevent misdiagnosis and help doctors treat correctly. Diagnosis imaging has a big role in detecting and understanding nasolabial cysts. CT scans and MRIs are the main tools doctors use in clinics nowadays to diagnose this case. A CT scan is good for knowing the real size and where exactly the cyst is, also shows if bones near the cyst have changed or been damaged because the cyst has been pressing long time [5]. But MRI is better for soft tissue imaging, and it is good when you want to see how the cyst is touching the other tissues around. Also, it helps to prove the lesion is not an odontogenic one, which helps again in diagnosis [6]. Even now, when better tools for diagnosis and surgery are there, the knowledge in books and articles about nasolabial cyst is still limited. Most time, it is only some case reports or very few case series. Still, there is a big missing part in the study that tells us fully about what is seen in the radiology and histology of these cysts. We need more studies like that to better understand, develop diagnostic methods, and help give better treatment to people. So, this study wants to help by showing a few patients who have a nasolabial cyst. We mainly look at their scan results and tissue exam findings. When we combine this with what we see in real patient cases, the study wants to show a clearer picture of how this cyst behaves, how to find it, and how to manage it. This will help doctors to make diagnoses more correctly and choose better treatment plans for people who have it.

Despite nasolabial cysts being well-described as rare benign lesions, there is limited comprehensive evidence integrating their clinical presentation with radiological and histopathological features, as most existing literature is restricted to isolated case reports and small case series. This lack of consolidated data makes accurate diagnosis challenging and may lead to misclassification with other facial cysts. This study aimed to analyze the clinical, histopathological, and radiological features of nasolabial cysts and evaluate their association with symptom duration, cyst size, and bone remodeling.

**METHODS**

The purpose of this retrospective case series study was to examine the radiological and histological characteristics of nasolabial cysts. All patients involved in the study gave their informed consent, and the Ethical Consideration was

obtained from IRB # ATMC/IERC/13th (01-2023)/24 of Al-Tibri Medical College and Hospital, Isra University Karachi. The study comprised ten patients who received a nasolabial cyst diagnosis between January 24 and December 24. The following inclusion criteria were used to choose the patients: full radiological data availability, histopathological confirmation of nasolabial cysts, and clinical presentation of swelling in the nasolabial region. The study excluded patients with insufficient medical data and those who had previously undergone surgery for nasolabial cysts. The required sample size is 11 individuals. The patients' medical records were examined to gather information on their age, gender, and ethnicity as well as their clinical presentation, size, and location of cysts, length of symptoms, and modes of therapy. Pathology reports provided the histological data, while computed tomography (CT) scans provided the radiological data. Radiological Evaluation: The size, location, and extent of the cysts, together with any related bone remodeling or scalloping, were evaluated by reviewing the CT scans. Upon independent assessment of the CT scans by two radiologists, disagreements were settled by consensus. Histopathological Analysis: Cyst specimens taken following surgical excision were subjected to a histological investigation. The presence of mucin-secreting cells, pseudostratified columnar epithelium, and any indication of squamous metaplasia was evaluated in the cyst lining. Two pathologists examined each slide separately, and when they disagreed, they came to a consensus. The clinical and demographic features of the patients were compiled using descriptive statistics. The standard deviation (SD) of continuous data, such as the age and size of the cysts, was represented by the mean ± SD. Frequencies and percentages were used to represent categorical characteristics, such as gender, ethnicity, and histological results. The relationship between the length of symptoms and the existence of bone remodeling on CT scans was assessed using the Kruskal-Wallis test. Statistical significance was attained when the p-value was less than 0.05. Version 22.0 of the SPSS software was used for all statistical analyses.

**RESULTS**

The results of the study highlight several key findings related to nasolabial cysts. Descriptive statistics showed that the mean age of patients was 44.9 ± 12.1 years, with cysts averaging 2.8 ± 1.1 cm in size and a mean symptom duration of 3.7 ± 2.0 years (Table 1).

**Table 1:** Descriptive Statistics

Variables	Mean ± SD
Age (Years)	44.9 ± 12.1
Size of Cyst (cm)	2.8 ± 1.1

Duration of Symptoms (Years)	3.7 ± 2.0
------------------------------	-----------

Gender distribution was 60% male and 40% female, with a higher prevalence among the Sindhi ethnic group (50%). CT findings revealed that 50% of the patients had no bone changes, while 40% exhibited scalloping and 30% showed bone remodeling (Table 2).

**Table 2:** Frequency Distribution

Variables	Categories	Frequency (%)
Gender	Male	6 (60%)
	Female	4 (40%)
Ethnicity	Sindhi	5 (50%)
	Balochi	2 (20%)
	Urdu	3 (30%)
CT Findings	None	5 (50%)
	Scalloping	4 (40%)
	Remodeling	1 (10%)
Histopathology	Pseudostratified Columnar	4 (40%)
	Mucin-Secreting Columnar	3 (30%)
	Double-Layer Columnar and Cuboidal	1 (10%)
	Squamous Metaplasia	2 (20%)

A significant association was found between the duration of symptoms and CT findings ( $p=0.030$ ), with longer symptom durations linked to scalloping or remodeling (Table 3).

**Table 3:** Association analysis between CT Findings and Duration of Symptoms

CT Findings	Mean Duration of Symptoms (Years)	H-Value	p-Value
None	2.2 ± 0.8	8.21	0.016
Scalloping	4.0 ± 1.2		
Remodeling	5.5 ± 0.9		

Kruskal-Wallis test was applied,  $p$ -value < 0.050

However, there was no statistically significant difference in age between patients with and without CT changes ( $p=0.060$ ), suggesting that age does not strongly predict bone involvement (Table 4).

**Table 4:** Relationship between Age Difference Based on CT Findings

CT Findings	Mean Age of Symptoms (Years)	t-Value	p-Value
None	38.6 ± 11.0	2.13	0.060
Scalloping/Remodeling	51.3 ± 9.6		

An independent t-test was applied,  $p$ -value < 0.050

The correlation between cyst size and duration of symptoms was also significant ( $r=0.32$ ,  $p=0.009$ ), indicating that larger cysts were associated with longer symptom durations. Overall, the findings underscore the importance of early diagnosis and treatment to prevent complications like bone remodeling, particularly in patients with longer-standing cysts (Table 5).

**Table 5:** Correlation Between Cyst Size and Duration of Symptoms

Variable 1	Variable 2	Correlation Coefficient (r)	p-Value
Cyst Size (cm)	Duration of Symptoms (Years)	0.32	0.009

Pearson correlation coefficient  $p$ -value < 0.050

## DISCUSSION

This study provides a comprehensive analysis of the clinical, radiological, and histopathological features of nasolabial cysts, enhancing the existing understanding of this Rare but significant lesion. Nasolabial cysts are non-odontogenic developmental lesions arising from remnants of the embryonic nasolacrimal duct, typically presenting as a painless swelling in the nasolabial fold. In our cohort, the mean patient age was  $44.9 \pm 12.1$  years, with a slight male predominance (60%). These findings align with previously reported trends, where nasolabial cysts commonly affect middle-aged individuals, particularly in Asian populations [2, 7]. CT imaging revealed maxillary bone remodeling in 30% of patients, scalloping in 40%, and no significant changes in 50%. These findings are in line with prior studies that have shown imaging, particularly CT, to be critical for preoperative planning and for evaluating bone involvement [5, 8]. Study found a moderate correlation ( $r=0.32$ ,  $p=0.009$ ) between cyst size and symptom duration, suggesting that larger cysts tend to be present for longer periods, a trend also noted in recent case reviews [9, 10]. The mean cyst size was  $2.8 \pm 1.1$  cm, with an average symptom duration of  $3.7 \pm 2.0$  years. Almutairi *et al.*, highlighted that nasolabial cysts may remain asymptomatic for extended periods but can gradually enlarge and compromise aesthetics and function [2]. Though a t-test comparing bone change presence with age produced a  $p$ -value of 0.06 (not statistically significant), patients with scalloping or remodeling were on average older ( $51.3 \pm 9.6$  years) than those without such features ( $38.6 \pm 11.0$  years). This implies that while age may play a role, symptom duration and cyst behavior are more likely to influence bone alterations [11, 12]. Overall, the current findings reaffirm that nasolabial cysts generally occur in the middle age group, with a slow-growing course and late onset of symptoms, supporting previous clinical observations [13, 14]. These cysts often go unnoticed until they attain sufficient size to cause visible facial asymmetry or nasal obstruction. Long-standing lesions are more prone to result in osseous changes, which have been previously documented by Yang *et al.*, [11] and Sato *et al.*, [15]. Advanced imaging modalities also play a critical role. While CT offers high-resolution bone detail, MRI, particularly with fat suppression techniques, can better define cystic content and rule out differential diagnoses [16]. In current study, prolonged symptom duration showed a strong correlation with radiological changes—specifically bone scalloping and remodeling on CT scans ( $p=0.03$ ),

underscoring the importance of timely diagnosis and intervention, as emphasized in the systematic review by Sheikh *et al.*, [17]. Histopathologically, all specimens demonstrated a pseudostratified columnar epithelial lining. Mucin cells were observed in 30% of cases, while squamous metaplasia was present in 20%, a pattern that supports earlier findings indicating that chronicity and local inflammation may influence epithelial transformation [18, 19]. Where MRI is unavailable, high-resolution ultrasound is a viable adjunctive diagnostic tool [20]. Histological evaluation remains pivotal in diagnosis, as seen in our study and those by Sheikh *et al.*, and Ramos *et al.*, where mucus-secreting cells and squamous metaplasia were variably present [17, 21]. This epithelial heterogeneity is consistent with the observations made by Ramos *et al.*, and others, who found histologic variability based on the duration and infectious status of the cyst [21]. Early case reports, including Wesley *et al.*, also emphasized similar patterns of long symptom latency and the importance of histological confirmation [22]. Regarding treatment, both sublabial excision and transnasal endoscopic marsupialization remain effective options. Current findings support existing literature suggesting that surgical technique selection should be based on surgeon expertise, lesion accessibility, and patient preference, with low recurrence rates reported for both methods [9, 21]. Moreover, psychological and cosmetic implications, particularly with bilateral or recurrent cysts, highlight the need for prompt surgical intervention [15, 22]. This study is limited by its single-center design and relatively small sample size, which may restrict the generalizability of the findings. Additionally, the lack of long-term postoperative follow-up limits assessment of recurrence rates and long-term surgical outcomes. More studies in the future should look at how treatment affects in the long term, and also at using less invasive techniques to manage this cyst.

## CONCLUSIONS

It was concluded that even if nasolabial cysts are not very common, they still cause many difficulties in clinical practice. Especially when symptoms last for a long time, this study shows how important early diagnosis and management are to stop complications like bone changes. Findings also suggest that the best way to handle this cyst is using a team approach with imaging, histology, and clinical examination all together.

## Authors' Contribution

Conceptualization: VKK, AMN

Methodology: VKK, SMMR, SK, SMAR, SBJ

Formal analysis: VKK, SMAR, SBJ, AMN

Writing and Drafting: SMMR, SK, FI, AMN

Review and Editing: VKK, SMMR, SK, SMAR, SBJ, FI, AMN

All authors approved the final manuscript and take responsibility for the integrity of the work

## Conflicts of Interest

All the authors declare no conflict of interest.

## Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

## REFERENCES

- [1] Mane BS and Gavali RM. Our experience at a tertiary medical college: excision with sublabial approach versus marsupialization with transnasal endoscopic approach in patients with nasolabial cyst. *Indian Journal of Otolaryngology and Head and Neck Surgery*. 2024 Feb; 76(1): 676-82. doi: 10.1007/s12070-023-04248-5.
- [2] Almutairi A, Alaglan A, Alenezi M, Alanazy S, Al-Wutayd O. Nasolabial cyst: case report and review of management options. *BioMed Central Surgery*. 2020 Dec; 20: 1-6. doi: 10.1186/s12893-020-0677-3.
- [3] Zhang J, Wu X, Ma J. A new transnasal approach of Nd: YAG laser treating nasolabial cysts. *Lasers in Medical Science*. 2022 Mar; 37(2): 1321-4. doi: 10.1007/s10103-021-03394-y.
- [4] Teixeira HR, Lopes MB, Abreu AP, Rosa MH, Antunes L. Transnasal marsupialization of nasolabial cyst: a case report. *International Journal of Otorhinolaryngology Clinics*. 2022 Jun; 14(2): 69-70. doi: 10.5005/jp-journals-10003-1404.
- [5] Singh AK, Tandon S, Malhotra T, Gaur S, Raj A. Nasolabial cyst: our experience and review of literature. *Journal of Coastal Life Medicine*. 2022; 10(3): 458-461.
- [6] Lee DH, Yoon TM, Lee JK, Lim SC. Surgical treatment of nasolabial cysts in a single institute. *The Journal of Laryngology and Otology*. 2020 Feb; 134(2): 141-4. doi: 10.1017/S0022215119002111.
- [7] Rodrigues BT, Fischer A, Romañach MJ, de Andrade BA, de Almeida Freire N, Israel MS. Nasolabial cyst in an elderly patient: A case report. *Gerodontology*. 2021 Sep; 38(3): 317-20. doi: 10.1111/ger.12521.
- [8] Tsai HW, Chao WC, Lee YH, Tsai YT, Tsai MS, Lee YC. Sublabial excision versus transnasal endoscopic marsupialization for nasolabial cysts: A systematic

- review and meta-analysis. *Clinical Otolaryngology*. 2024 Jan; 49(1): 102-8. doi: 10.1111/coa.14105.
- [9] Grover M, Nirwan S, Nahar R, Soni S. Management of Nasoalveolar Cyst by Transnasal Endoscopic Marsupialization. *Bengal Journal of Otolaryngology and Head Neck Surgery*. 2023 Sep;3 1(1): 1-6. doi: 10.47210/bjohns.2023.v3i1i.946.
- [10] Tilaveridis I, Venetis G, Tatsis D, Kalaitidou I, Zouloumis L. The aid of microsurgical instruments in nasolabial cyst enucleation. A report of two cases with critical review of the therapeutic approach. *Journal of Surgical Case Reports*. 2023 Jan; 2023(1): rjad011. doi: 10.1093/jscr/rjad011.
- [11] Yang H, Sun D, Tang Y, Su Y. Combined endoscope and low-temperature plasma knife for bilateral nasal vestibular cyst with sinus tract formation. *European Archives of Oto-Rhino-Laryngology*. 2023 Jan; 280(1): 479-81. doi: 10.1007/s00405-022-07751-7
- [12] Karki S, Shrestha BL, Adhikari P, Thapa NM. Surgical management of nasolabial cyst by transnasal endoscopic marsupialization: a Nepalese perspective. *Journal of Nepal Health Research Council*. 2021;19(1): 108-111.
- [13] Adhikari P, Shrestha BL, Sinha BK. Nasolabial cyst: review of 18 cases. *Journal of the Nepal Medical Association*. 2020;58(224):353-356. doi: 10.31729/jnma.5114.
- [14] Bhardwaj K, Jaiswani G, Sinha G, Kumar A, Gautam R, Raina A et al. Rare Case of Non-Odontogenic Nasolabial Cyst. *Journal of Population Therapeutics and Clinical Pharmacology*. 2025; 32(2): 377-381. doi: 10.53555/e2r5vq08.
- [15] Sato M, Morita K, Kabasawa Y, Harada H. Bilateral nasolabial cysts: a case report. *Journal of Medical Case Reports*. 2016 Dec; 10: 1-5. doi: 10.1186/s13256-016-1024-2.
- [16] Lk SK, Sidharthan D, Stanly B, Mathew CM. Nasolabial cyst: a diagnostic dilemma. *Journal of Dentistry*. 2022 Mar; 23(1): 72.
- [17] Sheikh AB, Chin OY, Fang CH, Liu JK, Baredes S, Eloy JA. Nasolabial cysts: a systematic review of 311 cases. *The Laryngoscope*. 2016 Jan; 126(1): 60-6. doi: 10.1002/lary.25433.
- [18] Su CY, Chien CY, Hwang CF. A new transnasal approach to endoscopic marsupialization of the nasolabial cyst. *The Laryngoscope*. 1999 Jul; 109(7): 1116-8. doi: 10.1097/00005537-199907000-00020.
- [19] Boffano P, Gallesio C, Campisi P, Roccia F. Diagnosis and surgical treatment of a nasolabial cyst. *Journal of Craniofacial Surgery*. 2011 Sep; 22(5): 1946-8. doi: 10.1097/SCS.0b013e31822ea751.
- [20] Toribio Y and Roehrl MH. The nasolabial cyst: a nonodontogenic oral cyst related to nasolacrimal duct epithelium. *Archives of Pathology and Laboratory Medicine*. 2011 Nov; 135(11): 1499-503. doi: 10.5858/arpa.2010-0338-RS.
- [21] Ramos TC, Mesquita RA, Gomez RS, Castro WH. Transnasal approach to marsupialization of the nasolabial cyst: report of 2 cases. *Journal of Oral and Maxillofacial Surgery*. 2007 Jun; 65(6): 1241-3. doi: 10.1016/j.joms.2005.10.049.
- [22] Wesley RK, Scannell T, Nathan LE. Nasolabial cyst: presentation of a case with a review of the literature. *Journal of Oral and Maxillofacial Surgery*. 1984 Mar; 42(3): 188-92. doi: 10.1016/S0278-2391(84)80032-4.