



Original Article



Beyond the Gag Reflex: Understanding Soft Palate Variations for Better Dental Care

Khezran Qamar¹, Maryam Mumtaz³, Muhammad Hammad Rafique¹, Khalid Yaqub³, Ayema Nazif¹, Tayyaba Hamid^{1,2} and Sajid Naeem¹

¹Department of Prosthodontics, Lahore Medical and Dental College, Lahore, Pakistan

²AFDB Dental Clinic, Illionis, United States of America

³Department of Prosthodontics, Institute of Dentistry, Combined Military Hospital Lahore Medical College, Lahore, Pakistan

ARTICLE INFO

Keywords:

Frequency, Complete Dentures, Gagging, Gender, Partial Dentures, Soft Palate

How to Cite:

Qamar, K., Mumtaz, M., Rafique, M. H., Yaqub, K., Nazif, A., Hamid, T., & Naeem, S. (2025). Beyond the Gag Reflex: Understanding Soft Palate Variations for Better Dental Care: Understanding Soft Palate Variations for Better Dental Care. *Pakistan Journal of Health Sciences*, 6(7), 204-208. <https://doi.org/10.54393/pjhs.v6i7.3258>

***Corresponding Author:**

Khezran Qamar
Department of Prosthodontics, Lahore Medical and Dental College, Lahore, Pakistan
khezranqamar@gmail.com

Received Date: 12th June, 2025

Revised Date: 21st July, 2025

Acceptance Date: 28th July, 2025

Published Date: 31st July, 2025

ABSTRACT

The gag reflex is a defense process to protect the airway from irritants. In a few patients, exaggerated reflexes jeopardize the dental treatment and compromise the dental procedures.

Objectives: To evaluate the potential association of the severity of gag reflex with types of soft palate in patients attending to prosthodontic clinic and to find out the association within gender, age groups. **Methods:** A cross-sectional observational study included 250 patients of both genders who needed dentures, were selected. Gag reflex was checked by touching the anterior to posterior part of the maxilla and mandible with a dental mirror while taking impressions. Gag Reflex Index (GRI) was used for the assessment of gag response. The reaction of the patients was recorded on V grades. The types of soft palates were classified via house classification as I, II, and III. **Results:** The soft palate forms frequently observed were soft palate drop type I, 60.0% followed by type II, 31.6% and 8.0% patients had type III soft palate drop. The gag reflex grade V was 5.6%, grade IV was 4.8%, grade III was 8.4%, grade II was 22.0% and grade I was 59.2%. The association of gag reflex grades with gender and types of soft palate drop was found to be significant, 0.020 and <0.001. However, with different age groups, insignificant results were obtained, p 0.070. **Conclusions:** The study concluded that the severity of gagging is found more in female patients and patients with type II and III soft palatal anatomy. Dentists need to evaluate different soft palatal types, especially when dealing with female patients needing prosthodontic treatment.

INTRODUCTION

The pharyngeal reflex, also known as the gag reflex, is a complex involuntary phenomenon in human physiology. Though the gag reflex is a protective defense response to keep the airway clear of irritants, in a few patients the extreme response elicits and jeopardizes the dental treatment [1]. During dental procedures, especially fabricating complete and partial dentures in prosthodontics, the frequency of gagging is common. The prosthodontic procedures, such as steps of making dentures like try selection, impression taking, determining vertical and horizontal jaw relation, denture insertion, and removal, elicit gag reflex. There can be external and internal stimuli that trigger the gagging response. External

stimuli such as the sight of a mouth mirror, impression trays, taste, and smell of impression materials can trigger the gag reflex [2,3]. Intraoral stimuli, such as the soft palate anatomy, show hyposensitive and hypersensitive responses. The trigger points of the gag reflex are associated with the types of soft palate. The soft palate is a muscular, mobile fold that is suspended from the hard palate posteriorly between the oropharynx and nasopharynx [4]. In the two domes of the hard palate, the muscles of the soft palate are inserted, and they hang down posteriorly and help in speech and swallowing [5]. There are three types of soft palate based on the angle that it forms with the hard palate. Type I is the soft palate angle



with the hard palate that is horizontal, shallow <10 degrees angle, and flat. Type II soft palate makes an angle between 10–45 degrees, whereas type III palate runs down, making a sharp angle of 70 degrees [6]. In denture patients, the posterior denture seal is made, and the posterior extension of the maxillary denture is always kept anterior to the mobile tissues of the soft palate. Any extension ahead of the movable soft palate tissues elicits gagging [7]. Patients with type II and III soft and hard palate anatomy greatly affect the posterior denture seal, as there is more mobile tissue and contact, and a lack of contact with the denture during speech and swallowing triggers gagging [8]. Prosthodontic treatment modalities to minimize the gagging reflex should be practiced. To cope with the gagging reflex and for practicing smooth dentistry, various prosthodontic modalities to control the gag reflex can be used [9]. The methods to manage gag reflex include psychological intervention, relaxation, distraction, desensitization, and cognitive behavior therapy. Practicing these modalities not only prevents disturbance, delay, and repetition of treatment but also saves time and cost of the procedure [10]. It is empirical to identify the severity of gagging in prosthodontic patients so that the dentist has the idea of dealing with the patient via standard treatment procedures or practicing alternative methods to control the gag reflex and get a successful and comfortable treatment outcome [11]. Care is needed, especially when dealing with geriatric patients during impression taking with alginate and impression compound [12]. Depending on the patient's condition and accurate diagnosis, the gagging can be controlled, and any disturbance that can jeopardize the steps of dental procedures can be rectified [11]. The soft palate plays a critical role in the gag reflex, and its morphology can influence the severity of gagging [13]. By understanding individual variations in gag reflex and soft palate anatomy, prosthodontists can tailor their approaches to minimize discomfort and anxiety and can improve patient compliance and treatment success rates. To the best of our knowledge, no study in our region has been conducted to evaluate the types of soft palate and gag reflex frequency in prosthodontic patients.

The gag reflex is a common challenge in prosthodontic practice, particularly during impression-making and denture fabrication. Although previous studies have explored gag reflex prevalence and management strategies, limited research has evaluated its association with specific soft palate morphologies using standardized classifications. Furthermore, regional data examining the relationship between gag reflex severity, soft palate types, gender, and age groups remain scarce. This highlights the need for locally generated evidence to better guide clinical assessment and treatment planning in prosthodontic patients. The study aimed to evaluate the potential

association of the severity of gag reflex with types of soft palate in patients attending to prosthodontic clinic and also find out the association within gender, age groups.

METHODS

A cross-sectional observational study of 3 months duration from 20th Feb 2025 till 20th May 2025 was carried out in the department of Prosthodontics of Lahore Medical and Dental College Lahore. Ethical clearance was obtained from the Ethical Review Board of the Dental College. Every patient was comfortably seated in the dental chair in upright position with straight legs on the chair and facing the dentist. A total of 250 patients of both gender; needing complete and partial dentures presented to the OPD of clinic with in the age range from 20 till 80 years were included in the study. The sample size was calculated from the RAO Soft Epi calculator against a 20.0% prevalence of type III gag reflex in patients, with 95% confidence interval, keeping margin of error 5%. Non-probability purposive sampling technique was used for sample selection. Healthy patients requiring complete and partial dentures within the selected age range were included. Patients with Congenital and developmental soft and hard tissue defects, palatopharyngeal incompetencies, a history of oral/maxillofacial trauma, and patients wearing obturators were excluded from study. Verbal consent was taken. The gag severity index (GSI) was used to find out the severity of gagging and the patients were divided into 5 grades based on the severity of gagging reflex. Grade V: Patients start to gag just by sitting into the dental chair on their first dental visit. [14] Grade IV: Patients start to gag during visual examination. Grade III: Patients who gag when the maxilla is touched by the mouth mirror from anterior palate to the post dam area and in the mandible when the mirror was moved from the labial sulcus to the lingual sulcus and to the reteromolar pad. Grade II: Patients who gag while taking primary impression with alginate and impression compound in stock trays. Grade I: Patients who felt little controllable gag during primary impression taking. Two experienced prosthodontists evaluated the gag reflex and evaluated the soft palate types and decided them by agreement. The types of soft palates noted were classified based on House Classification as I, II, III [15]. Class I: drop of soft palate concerning hard palate is horizontal, flat extension posteriorly, minimum muscle activity, making an angle with hard palate <10 degrees. Type II: The drop of soft palate is more than 45 degrees with a gradual slope and moderate muscle activity, making an angle with the hard palate from 10–45 degrees. Type III: The drop is more than a 45-degree angle concerning the hard palate, steep descent of the soft palate. The patients were divided into 3 groups concerning age, i.e., Group I: 20–40, Group II: 41–60 years, Group III: 61–80 years. The Data was analyzed using SPSS

Version 26. Qualitative variables were presented as frequencies and percentages, whereas the quantitative variables were presented as means and SD. Chi-square square used to find out the association of gag reflex with gender, age groups, and soft palate types. p-value <0.050 will be the significance level.

RESULTS

Out of 250 patients 111(44.4 %) were males and 139(55.6 %) were female patients. The patients age ranged from 20 to 80 years with mean age of 53.79±13.50 SD. 41 (16.4 %) of patients were of younger age group from 20–40 years, 120 (48.0 %) were from 41–60 years, and 88 (35.2%) patients presented were from older age group. The type of soft palate forms frequently observed was soft palate drop type I 151(60.0%) followed by type II 79(31.6%) and only 20(8.0 %) patients had type III soft palate drop. The gag reflex grade V was 14(5.6%), grade IV was 12(4.8%), grade III was 21(8.4%), grade II was 55(22.0 %), and grade I was 148 (59.2%)(Table 1).

Table 1: Frequency Distribution of Gag Reflex Grades and Types of Soft Palate Drops. N=250.

Gag Reflex Grades					Soft Palate Types		
Grade I n %	Grade II n %	Grade III n %	Grade IV n %	Grade V n %	Drop I n %	Drop II n %	Drop III n %
148 (59.2)	55 (22.0)	21(8.4)	12(4.8)	14 (5.6)	151 (60.0%)	79 (31.6)	20 (8.0)

The association of gag reflex grades with gender was found to be significant 0.020. The association of different age groups with gag reflex was found to be insignificant p 0.730 however, gag reflex severity (grade I-V) increases with the increasing grade of types of soft palate drop (I, II, III), and a significant association was found p value <0.001 (Table 2).

Table 2: Association of Gag Reflex Grades with Age Groups, Gender, and Soft Palate Types; N=250

Variables	Gag Reflex Grades					p-Value
	Grade I n (%)	Grade II n (%)	Grade III n (%)	Grade IV n (%)	Grade V n (%)	
Age Groups						
20–40 yrs	27 (65.9)	10 (24.4)	1 (2.4)	2 (4.9)	1 (2.4)	0.070
41–60 yrs	80 (66.7)	23 (19.2)	10 (8.3)	5 (4.2)	2 (1.7)	
61–80 yrs	55 (62.5)	21 (23.9)	10 (11.4)	0 (0.0)	2 (2.3)	
Gender						
Male	78 (52.7)	20 (36.4)	7 (33.3)	3 (25.0)	3 (21.4)	0.020
Female	70 (47.3)	35 (63.6)	14 (66.7)	9 (75.0)	11 (78.6)	
Soft Palate Types						
Drop I	129 (87.2)	18 (32.7)	2 (9.5)	1 (8.3)	1 (7.1)	<0.001
Drop II	18 (12.2)	35 (63.6)	17 (81.0)	8 (66.7)	1 (7.1)	
Drop III	1 (0.7)	2 (3.6)	2 (9.5)	3 (25.0)	12 (85.7)	

Significance level p<0.050

DISCUSSION

Dental practitioners frequently encounter the challenge of gagging during procedures, which can hinder or even prevent effective treatment despite advancements in modern dentistry. Understanding the causes and impact of gagging is essential for improving patient comfort and acceptance of dental treatment [16, 17]. In the present study, grade I gag reflex was the most frequent (59.2%), followed by grade II (22.0%), while severe grades IV (4.8%) and V (5.6%) were the least common. These findings align with those of Alamgir *et al.*, who reported grade I (40.0%) and grade II (32.7%) as the most frequent, with lower rates for grades IV (6.7%) and V (0.7%) [14]. Similarly, Meshni *et al.* found a 49.1% prevalence of gagging in their patients [18]. Current results showed a higher proportion of female patients (55.6%) compared to males (44.4%). Although grade I was the most common in both genders, severe gag reflexes were more frequent in females. Specifically, grade IV occurred in 75.0% of females and 25.0% of males, while grade V was observed in 78.6% of females and 21.4% of males. The association between gag reflex grades and gender was statistically significant (p=0.02). These findings agree with Alamgir *et al.*, who also reported a higher prevalence of exaggerated gag reflex in females (59.3%) compared to males (40.7%) [14], as well as studies from Saudi Arabia and other regions reporting similar trends [19–21]. Gender differences in jaw size and psychological sensitivity, as suggested by Stefos *et al.*, may partially explain these findings [22]. Regarding soft palate morphology, type I was the most common (60.0%), followed by type II (31.6%) and type III (8.0%). The association between gag reflex severity and soft palate type was highly significant (p<0.001). Patients with type I soft palate predominantly exhibited grade I gag reflex (87.2%), whereas severe grades IV and V were rare (8.3% and 7.1%, respectively). In contrast, type II was more frequently associated with higher gag reflex grades, including grade IV (66.7%). Type III showed the most severe response, with 85.7% of patients having grade V gag reflex. These observations correspond with Halboub *et al.*, who reported a significant correlation between soft palate type and gagging, with more severe gagging in type II and III morphologies [19]. Anatomically, a steep drop from the hard palate and reduced posterior surface area in types II and III may predispose to increased gag sensitivity due to reduced tolerance for impression trays [4]. Age did not show a statistically significant association with gag reflex severity (p=0.730, not significant). However, most patients were middle-aged (41–60 years; 48.0%), followed by older adults (61–80 years; 35.2%), with younger adults (20–40 years) representing the smallest group (16.4%). Grade I gag reflex was the most frequent across all age groups, while

severe grades IV and V were uncommon. Similar findings were reported by Alamgir et al., where a mild gag reflex was more prevalent in younger patients, while a moderate gag reflex appeared more in middle-aged individuals [14]. The study was conducted in a single clinic, limiting generalizability. Its cross-sectional design prevents causal conclusions. Psychological factors affecting the gag reflex were not evaluated, and subjective grading may introduce bias.

This study was limited by its single-center, cross-sectional design, which restricts causal inference and generalizability. The use of purposive sampling and subjective clinical grading may introduce observer bias, and psychological factors influencing gag reflex were not assessed. Future multicenter studies with larger, more diverse populations and incorporation of objective assessment tools and psychological profiling are recommended. Longitudinal research may further clarify the predictive value of soft palate morphology in managing gag reflex during prosthodontic treatment.

CONCLUSIONS

The severity of gagging is found more in female patients and patients with type II and III soft palatal anatomy. Dentists need to evaluate different soft palatal types, especially when dealing with female patients needing prosthodontic treatment.

Authors' Contribution

Conceptualization: TH

Methodology: KQ, AN, TH

Formal analysis: MHR

Writing and Drafting: MM, KY, SN

Review and Editing: MM, KY, SN, MHR, KQ, AN, TH

All authors approved the final manuscript and take responsibility for the integrity of the work

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Mehdizadeh M, Mohammadbeigi A, Sharifinejad A. An Overview About New Methods in Management of Gag Reflex During Dental Treatment: A Systematic Review. *Journal of Dentistry*.2023Dec;24(4):372. doi: 10.30476/dentj.2023.98310.2164.
- [2] Meenakshi S, Dureja S, Kavita GC, Pallavi M, Swamy KN, Kottur A, et al. The Gag Reflex: A Hurdle in Dentistry-Literature Review. *Journal of Pharmaceutical Research International*. 2021 Oct; 33(46B): 224-37.doi:10.9734/jpri/2021/v33i46B32937.
- [3] Hassan H, Rafique A, Andleeb S, Javaid M, Ahmad H, Khalid S. Post-Insertion Complaints Associated with Complete Denture Prosthesis in a Tertiary Care Hospital. *Rawal Medical Journal*.2023 Mar; 48(1): 200.
- [4] Mortazavi H, Hazrati P, Azadi A. The Main Differential Diagnosis of Swellings in the Posterolateral Part of the Hard Palate: A Mini Review. *Journal of Iranian Medical Council*.2023Aug;6(4).doi:10.18502/jimc.v6i4.13439.
- [5] Prasad V, Chhodon L, Aravindhan A, Zaidi SJ. Prosthetic Rehabilitation of Patients with Cleft Lip and Palate. *Journal of Cleft Lip Palate and Craniofacial Anomalies*.2022Jul;9(2):189-96.doi: 10.4103/jclpca.jclpca_6_22.
- [6] Yuan Z, Yuan J, Wang Z, Rong Q. Rehabilitation of Combined Hard and Soft Palate Defects with a Soft Relining Prosthesis: A Clinical Report. *The Cleft Palate Craniofacial Journal*.2024Aug;10556656241272481. doi: 10.1177/10556656241272481.
- [7] Qamar K, Rizvi SH, Yasser F, Aamir H, Ahmed RA, Muzaffar M, et al. Patient Satisfaction with Quality of Dentures at Teaching Hospital of Lahore: Quality Dentures and Patient Satisfaction. *Pakistan Journal of Health Sciences*.2024May;5(05):25-9.doi:10.54393/pjhs.v5i05.1451.
- [8] Hassan H, Rafique A, Andleeb S, Javaid M, Ahmad H, Khalid S. Post-Insertion Complaints Associated with Complete Denture Prosthesis in a Tertiary Care Hospital. *Rawal Medical Journal*.2023 Mar; 48(1): 200.
- [9] Colvenkar S, Reddy V, Thotapalli S, Rani KD, Bharadwaj S, Ali S, et al. A Simple Step-by-Step Technique for the Management of Gagging in Edentulous Patient. *Cureus*.2022Apr;14(4).doi: 10.7759/cureus.24423.
- [10] Ali S, George B, Kirmani U, Al-Saiari AK, Almasabi FR, Iqbal Z. Gagging and Its Management in Prosthodontic Patients: A Review of Literature. *Biomedica*. 2018 Jul; 34(3): 179.
- [11] Mehdizadeh M, Mohammadbeigi A, Sharifinejad A. An Overview About New Methods in Management of Gag Reflex During Dental Treatment: A Systematic Review. *Journal of Dentistry*.2023Dec;24(4):372. doi: 10.30476/dentj.2023.98310.2164.
- [12] Gali S, Hithyshi GK. Managing a Hyper Gag Reflex Edentulous Patient with Modeling Compound and Training Dentures. *Manipal Journal of Dental Sciences*. 2020; 5(1): 35-8.
- [13] Parakh MK. Gagging and Its Management-A Review. *Oral Science International*.2022 Sep; 19(3): 141-6. doi: 10.1002/osi2.1128.

- [14] Alamgir K, Saleem T. Prevalence and Severity of Gag Reflex in Patients Presenting at the Department of Prosthodontics. *The Professional Medical Journal*. 2024 Sep; 31(09):1301-6. doi:10.29309/TPMJ/2024.31.09.8087.
- [15] Chaturvedi S, Khaled Addas M, Al Humaidi AS, Al Qahtani AM, Al Qahtani MD. A Novel Approach to Determine the Prevalence of Type of Soft Palate Using Digital Intraoral Impression. *International Journal of Dentistry*. 2017; 2017(1):3268064. doi:10.1155/2017/3268064.
- [16] Karademir Göceroğlu AH, Ölmez A. Prevalence of Gagging in Children and the Association Between Gagging and Dental Fear: A Cross-Sectional Study. *Türkiye Klinikleri Journal of Dental Sciences*. 2022 Jul; 28(3). doi: 10.5336/dentalsci.2021-87792.
- [17] Manoharan S, Subhashree R, Humar MS. Gag Reflex and Prosthodontic Procedures—A Comparative Study. *International Journal of Dentistry and Oral Sciences*. 2021 Mar; 8(03):1829-31. doi:10.19070/23 77-8075-21000363.
- [18] Meshni AA. Gag Reflex: A Comparative Study Among Different Prosthodontic Treatment Modalities. *Oral Health and Dental Management*. 2017; 16(4):1-4.
- [19] Halboub E, Sayed BA, Jaafari SA, Abutaleb GK, Arishi TM, Khadhi AH, et al. The Possible Association Between the Types of Soft Palate and Gag Reflex: A Preliminary Study. *Brazilian Dental Science*. 2023 Mar; 26(2). doi:10.4322/bds.2023.e3628
- [20] Kassab NH, Al-Saffar M. Gagging: A Problem in Prosthetic Dentistry and Its Medical Treatment. *Al-Rafidain Dental Journal*. 2005 Dec; 5(2):168-73. doi: 10.33899/rden.2005.45498.
- [21] Gucyetmez Topal B, Falay Civelek SB, Tiras M, Yigit T. The Prevalence and Influencing Factors of Gag Reflex in Children Aged 7-14 Years in the Dental Setting. *Journal of Oral Rehabilitation*. 2023 May; 50(5): 376-82. doi: 10.1111/joor.13432.
- [22] Stefos S, Zoidis P, Nimmo A. Managing Gag Reflex During Removable Partial Denture Treatment: A Review and a Clinical Report. *Journal of Prosthodontics*. 2019 Jul; 28(6): 618. doi: 10.1111/jopr.12957.